

Baza pytań do Lekarskiego Egzaminu Końcowego
Obstetrics and gynecology

Pytanie nr 1

Symptoms of ovarian cancer **do not** include:

- A. pain in the lower abdomen or pelvis.
- B. oliguria.
- C. increase in the abdominal circumference.
- D. tympanites, abdominal distention.
- E. urge incontinence.

Pytanie nr 2

Which statement regarding the placenta is **false**?

- A. forms the afterbirth together with the umbilical cord and the amniotic sac.
- B. its mass is approximately 500g.
- C. fully formed placenta is not observed until the second half of the second trimester.
- D. cotyledon is the morphological and structural unit of the placenta.
- E. right umbilical vein undergoes atrophy in the early stage of foetal development.

Pytanie nr 3

Which of the following are regarded as anatomical factors which can cause a miscarriage?

- A. endometriosis in the lesser pelvis.
- B. congenital defects of the uterus.
- C. cervical incompetence.

D. uterine fibroids.

E. all the above.

Pytanie nr 4

Which stage of endometriosis according to the classification of the American Society for Reproductive Medicine should be diagnosed in a patient with an endometrial cyst of the left ovary of 4 cm in diameter, partial obliteration of the recto-uterine pouch and superficial lesions of the peritoneal membrane of the diameter < 1 cm?

A. I.

B. II.

C. III.

D. IV.

E. V.

Pytanie nr 5

The most common complication in twin monoamniotic pregnancies is:

A. preterm birth.

B. umbilical cord entanglement.

C. intrauterine growth restriction.

D. polyhydramnios.

E. umbilical cord prolapse.

Pytanie nr 6

Which of the following decreases men's fertility by lowering testosterone production?

A. ethanol.

B. ketoconazole.

C. spironolactone.

D. sulfasalazine.

E. all the above.

Pytanie nr 7

The range of normal values of Amniotic Fluid Index between 5 and 25 cm may be applied to the patients:

A. during the whole singleton pregnancy.

B. independently of the number of fetuses.

C. in singleton pregnancy only after 20 weeks of gestation.

D. in singleton pregnancy only after 24 weeks of gestation.

E. in singleton pregnancy only after 32 weeks of gestation.

Pytanie nr 8

Which of the following **is not** an indication for cesarean section?

A. Marfan syndrome with aorta dilatation and the coarctation of the aorta that has not been surgically corrected.

B. anxiety of a pregnant woman before the delivery.

C. significant difference of foetal weights of > 20% in twin pregnancy.

D. foetal weight > 3500 g in the longitudinal foetus breech position.

E. heart failure or significant arrhythmia in a pregnant woman.

Pytanie nr 9

A neglected transverse lie is characterized with:

- 1) impaction of the presenting shoulder in the pelvic inlet;
- 2) premature rupture of the membranes;
- 3) total cervical effacement;
- 4) excessive dilation of the lower uterine wall;
- 5) contraction of the uterus on a highly bent fetus.

The correct answer is:

- A. 1,2,3.
- B. 1,3,5.
- C. 2,4.
- D. 3,4,5.
- E. all the above.

Pytanie nr 10

The incidence of which of the following diseases **does not** increase in female patients with Turner syndrome?

- A. strabismus.
- B. grass pollen allergy.
- C. chronic otitis media.
- D. arterial hypertension.
- E. autoimmune thyroid disease.

Pytanie nr 11

Which of the following are regarded as risk factors for a transverse lie?

- 1) oligohydramnios;
- 2) polyhydramnios;
- 3) placenta praevia;
- 4) multiple pregnancy;
- 5) single pregnancy.

The correct answer is:

- A. 1,2,3,4.
- B. 1,3.
- C. 1,2,4,5.
- D. 1,3,5.
- E. 2,3,4.

Pytanie nr 12

A 35-year-old patient came to the gynecological outpatient clinic because of vaginal bleeding occurring between periods for about a year. The bleeding with dark brown blood occurs on 11th-12th day of the cycle and lasts 2-3 days. The periods occur every 30 days. The patient underwent a caesarian section two years earlier. Which of the following is the most probable cause of the bleedings?

- A. undiagnosed disorder of the clotting system.
- B. corpus luteum insufficiency.
- C. pelvic inflammatory disease.
- D. periovulatory LH surge.
- E. scar defect of the myometrium after the cesarean section.

Pytanie nr 13

A female patient was diagnosed with a lack of ovulation, acne and hirsutism. Those symptoms may suggest polycystic ovary syndrome. However, a presence of striae, a buffalo hump and arterial hypertension may suggest the diagnosis of Cushing's syndrome. Which of the following should be performed in order to differentiate those disorders?

- A. estradiol level.
- B. dexamethasone suppression test.
- C. metoclopramide test.
- D. lutropin level.
- E. follitropin level.

Pytanie nr 14

Which class in the Quigley scale corresponds to complete androgen insensitivity syndrome (CAIS)?

- A. from 1 to 3.
- B. 4.

C. 7.

D. 6.

E. 5.

Pytanie nr 15

Indicate the true statements concerning labor induction:

- 1) in the case of cholestasis of pregnancy when fatty acid concentration is 45 nmol/L the labor induction should be considered after 34 weeks of gestation;
- 2) in the case of gestational diabetes the labor induction should be considered after 39 weeks of gestation;
- 3) rupture of membranes is an absolute contraindication to labor preinduction with the use of prostaglandins;
- 4) in the case of pregestational diabetes the labor induction should be considered after 39 weeks of gestation;
- 5) in the pregnancy complicated by diabetes the labor induction is indicated when the difference between the fetal abdomen diameter and the biparietal diameter is over 4 cm.

The correct answer is:

A. 1,2,5.

B. 1,2,3.

C. 1,2,3,5.

D. only 2.

E. 3,4.

Pytanie nr 16

The indications for elective cesarean section in twoen pregnancy **do not** include:

A. each situation of abnormal presentation of any of the foetuses.

B. death of the foetus II.

C. TTTs syndrome (twin-to-twin syndrome).

D. significant difference in foetuses weight > 20%.

E. non-cephalic presentation of foetus I.

Pytanie nr 17

Which of the following surgical approaches is characterized by invisible scar, better healing in obese women but the limited field of view?

- A.** abdominal approach - the incision in the midline.
- B.** abdominal approach - the incision in the paramedian line.
- C.** abdominal approach - the Pfannenstiel incision.
- D.** vaginal approach.
- E.** laparoscopic approach.

Pytanie nr 18

Medical history of cervical incompetence is an indication for:

- A.** prophylactic placement of a cervical cerclage.
- B.** placement of a cervical cerclage when uterine contractions occur before 28th week of gestation.
- C.** placement of a cervical cerclage when there are symptoms of vaginal infection, along with antibiotic therapy.
- D.** frequent monitoring of the cervical length with ultrasonography.
- E.** calcium-rich diet in order to improve the consistence of the cervix.

Pytanie nr 19

An absolute indication for primary cesarean section in breech presentation **is not**:

- A.** foetus weight above 3500 g.
- B.** primipara.
- C.** umbilical cord prolapse.
- D.** placenta previa.

E. placenta abruption.

Pytanie nr 20

Which of the following determinations is used to diagnose hyperandrogenism in a 25-year-old woman?

A. ratio of LH to FSH.

B. free testosterone or index of free androgens.

C. dihydrotestosterone.

D. SHBG.

E. SHBG and dihydrotestosterone.

Pytanie nr 21

Untreated iron-deficiency anemia in a pregnant woman may lead to:

1) intrauterine growth retardation;

2) premature labour;

3) elongation of the pregnancy;

4) congenital anomalies;

5) prolonged labour.

The correct answer is:

A. 1,2,4,5.

B. 1,3,4,5.

C. 1,2,3,4.

D. 1,2,3,5.

E. 3,4,5.

Pytanie nr 22

The high risk factors of gestational diabetes include:

1) obesity of the pregnant woman (BMI > 30);

2) diabetes in the first-degree relatives;

3) birth weight of neonates of previous pregnancies > 4000 g;

4) age of the pregnant woman < 25 years;

5) no significant obstetric history.

The correct answer is:

- A. 1,2,3.
- B. 1,2,5.
- C. 2,3,4.
- D. 1,3,4.
- E. all the above.

Pytanie nr **23**

Which of the following **is not** typical of cervical cancer?

- A. urinary urgency.
- B. bloody vaginal discharge with unpleasant smell.
- C. intermenstrual bleeding.
- D. abdominal pain.
- E. contact bleeding.

Pytanie nr **24**

Which of the following **is not** among the abnormalities of the head position during labour:

- A. deep transverse arrest.
- B. brow presentation.
- C. occiput posterior position.
- D. left or right posterior position.
- E. occiput anterior position.

Pytanie nr **25**

When, during pregnancy, the concentration of the β -subunit of hCG in blood is the highest?

- A.** 5-6 week.
- B.** 6-7 week.
- C.** 9-12 week.
- D.** 14-16 week.
- E.** 16-18 week.

Pytanie nr 26

Which of the following is characteristic of borderline ovarian tumors?

- A.** no destructive infiltration of the stroma.
- B.** occurrence limited to the ovaries only.
- C.** prognosis similar to that of ovarian cancer.
- D.** no effective treatment options available.
- E.** occurrence in postmenopausal women only.

Pytanie nr 27

The administration of specific immunoglobulin is mandatory in the case of the pregnant, who had not been affected or vaccinated, and who had contact with a patient infected with:

- 1) type B hepatitis virus;
- 2) CMV virus;
- 3) fifth disease virus;
- 4) rubella virus;
- 5) chickenpox virus.

The correct answer is:

- A.** 1,3,5.
- B.** 2,3,5.
- C.** 2,3,4.
- D.** 1,4,5.

E. 4,5.

Pytanie nr 28

Microscopic evaluation of the material obtained at the outer opening of the cervix is called cervical smear (Papanicolaou test) and is used to diagnose precancerous states or cervical cancer. Which of the following should be advised in the case of the abnormal result of cervical smear - HSIL (*high-grade squamous intraepithelial lesion*)?

- A. repeat cervical smear - twice every 4-6 months.
- B. perform colposcopy with the biopsy of the suspected lesions and abrasion of the cervical canal.
- C. HPV determination.
- D. repeat cervical smear and determine HPV.
- E. none of the above.

Pytanie nr 29

A 64-year-old obese female reports postmenopausal vaginal bleeding. A cytological smear was obtained and in a transvaginal ultrasound endometrial hyperplasia was found. The mucosa biopsy was performed and the histopathological diagnosis of adenocarcinoma was obtained. Pap smear was normal. The uterus with appendages was completely resected and the lymph nodes were sampled. In postoperative examination adenocarcinoma cells were found in the uterus region (no evidence of serous membrane infiltration) and in the cervical canal near the internal orifice. No evidence of adnexal infiltration or metastases to the lymph nodes was found. Indicate tumor advancement according to FIGO classification:

- A. IA.
- B. IB.
- C. II.
- D. IIIA.
- E. IIIB.

Pytanie nr 30

The possible causes of cervical incompetence include:

- 1) inherited disorders;
- 2) traumas of the cervix;
- 3) state after the cervical conization;
- 4) multiple pregnancy.

The correct answer is:

- A.** only 1.
- B.** 1,2.
- C.** only 4.
- D.** all the above.
- E.** none of the above.

Pytanie nr 31

A 25-year-old man with normal male phenotype is examined because of infertility. Apart from a decreased sperm count in the semen analysis, gynecomastia was found and laboratory tests showed decreased follicle-stimulating hormone levels and increased testosterone levels. The most probable cause of the above abnormalities is:

- A.** chemotherapy in the past.
- B.** radiotherapy in the past.
- C.** use of anabolic steroids.
- D.** Kallmann syndrome.
- E.** Sertoli cell-only syndrome.

Pytanie nr 32

What percentage of the patients with diagnosed ovarian cancer are BRCA1/BRCA2 germinal mutation carriers?

- A.** 1%.
- B.** 5%.
- C.** 10%.

D. 30%.

E. 50%.

Pytanie nr 33

In a pregnant vegetarian you should expect the insufficiency of vitamin:

- 1) B₂;
- 2) B₁₂;
- 3) D;
- 4) C;
- 5) E.

The correct answer is:

A. only 1.

B. only 2.

C. only 3.

D. 1,3,4,5.

E. 1,2,4,5.

Pytanie nr 34

The normal age of menarche in Poland is defined as:

A. 8-9 years.

B. 9-16 years.

C. 16-17 years.

D. >18 years.

E. no norm exists.

Pytanie nr 35

Which of the following **cannot be** a consequence of intrauterine infection?

A. Down syndrome.

- B.** hydrocephalus.
- C.** fetal tachycardia.
- D.** preterm labor.
- E.** prelabor rupture of membranes.

Pytanie nr **36**

Which of the following concerning hyperprolactinemia **is false**?

- A.** sleep and stress are physiological conditions associated with increased secretion of prolactin.
- B.** hyperprolactinemia is the cause of 30% of cases of secondary amenorrhea after discontinuation of oral contraceptives.
- C.** drugs used in hyperprolactinemia treatment are cabergoline and bromocriptine.
- D.** consequences of hyperprolactinemia include the lack of pubic hair and a decrease in the secretion of adrenal androgens.
- E.** galactorrhoea is found in 30-90% women with hyperprolactinemia.

Pytanie nr **37**

A 33-year-old pregnant woman in 37th week of her second pregnancy came by the obstetric out-patient clinic because of the weakening of fetal movements. The course of the previous pregnancy was uncomplicated. The cardiotocography showed normal reactive FHR. Ultrasound examination of biophysical profile of the fetus revealed normal volume of amniotic fluid, one episode of breathing movements lasting 60 seconds, three movements of the trunk and twice hand closing and opening. The result of biophysical profile is:

- A.** 2/10 points.
- B.** 4/10 points.
- C.** 6/10 points.
- D.** 8/10 points.
- E.** 10/10 points.

Pytanie nr 38

The indications for insemination include:

- 1) decreased parameters of the semen;
- 2) ejaculation disorders;
- 3) PCOS;
- 4) III grade endometriosis;
- 5) idiopathic infertility.

The correct answer is:

- A.** 1,2,3.
- B.** 1,2,3,4.
- C.** 1,2,4,5.
- D.** 1,2,5.
- E.** 3,4,5.

Pytanie nr 39

If hypogastric pain appears approx. 2-3 days after the administration of methotrexate in a patient treated for ectopic pregnancy, you should:

- A.** perform urgent laparotomy with the longitudinal incision - this symptom means that there is haemorrhage because of ectopic pregnancy rupture.
- B.** perform urgent laparoscopy - this symptom means that there is haemorrhage because of ectopic pregnancy rupture.
- C.** perform urgent laparotomy with the transverse incision - this symptom means that there is haemorrhage because of ectopic pregnancy rupture.
- D.** observe the patient and monitor decrease in the hCG level - this is a typical pain after the treatment with methotrexate.
- E.** discharge the patient and inform her that the pain is most probably caused by ovulation.

Pytanie nr 40

In a patient with Turner syndrome (karyotype 45,X), within sexual organs, one can observe:

- A.** aplastic ovaries.
- B.** additional ovaries.
- C.** testes and ovaries.
- D.** testes only.
- E.** normal ovaries.

Pytanie nr 41

In women using biphasic hormonal contraception a typical menstrual period:

- A.** is short and light.
- B.** does not occur.
- C.** occurs irregularly and last for many days.
- D.** is heavy and long.
- E.** occurs every 21 days and is light.

Pytanie nr 42

Indicate the true statement concerning the uterine artery:

- A.** originates from the ovarian artery.
- B.** gives the branch running through the round ligament.
- C.** originates from the internal iliac artery.
- D.** crosses the ureter from the rear.
- E.** provides blood to the uterus only.

Pytanie nr 43

The prevention of Rh disease during pregnancy consists in the administration to each pregnant woman:

- A.** who is Rh-D negative, of 300 µg anti-RhD immunoglobulin at 28th-32th week of

gestation.

B. irrespectively of her Rh-D status, of 300 µg anti-RhD immunoglobulin at 28th-32th week of gestation.

C. who is Rh-D negative, of 300 µg anti-RhD immunoglobulin at 28th-32th week of gestation, but only if Rh-D status of the father is provided.

D. who is Rh-D negative, of 150 µg anti-RhD immunoglobulin at 28th-32th week of gestation.

E. who is Rh-D negative, of 300 µg anti-RhD immunoglobulin after 34th week of gestation.

Pytanie nr 44

Which of the following situations occurring during pregnancy requires a quick termination of the pregnancy?

A. premature placental abruption.

B. cervical cancer in pregnancy.

C. HIV infection.

D. adnexal torsion.

E. acute appendicitis.

Pytanie nr 45

Indicate the maneuver used in vaginal delivery of a baby in a longitudinal lie with breech presentation in order to release the child's shoulders:

A. Løvset's maneuver.

B. Mauriceau-Smellie-Veit maneuver.

C. Wiegand-Martin-von Winckel maneuver.

D. Gaskin maneuver.

E. Kiwisch-Scanzoni maneuver.

Pytanie nr 46

Which of the following is in favor of the diagnosis of antiphospholipid syndrome?

- A. vascular thrombosis.
- B. miscarriages.
- C. presence of anti-cardiolipin antibodies.
- D. presence of antibodies against β 2-glycoprotein 1.
- E. all the above.

Pytanie nr 47

Which of the following medications is certified for the treatment of stress urinary incontinence?

- A. paroxetine.
- B. duloxetine.
- C. venlafaxine.
- D. dapoxetine.
- E. fluoxetine.

Pytanie nr 48

Which of the following **is not** considered in the assessment of the cervix according to the Bishop score?

- A. dilation.
- B. consistency.
- C. distance from the sacral bone.
- D. effacement.
- E. fetal station.

Pytanie nr 49

Indicate the medications that can be safely used in pregnant patients:

- 1) aluminum compounds in the treatment of gastroesophageal reflux;
- 2) non-steroidal anti-inflammatory drugs from the group of non-selective cyclooxygenase inhibitors;
- 3) nystatin in the treatment of infectious diseases;
- 4) nitrofurantoin and furazidin in the treatment of urinary tract infections;
- 5) angiotensin converting enzyme inhibitors in the treatment of hypertension.

The correct answer is:

- A.** 1,4.
- B.** 1,2,4.
- C.** 2,3,5.
- D.** 1,3.
- E.** only 1.

Pytanie nr 50

Bacterial vaginitis:

- 1) is asymptomatic in most women, and in symptomatic cases manifests with a grey or white vaginal discharge;
- 2) is related to dysuria and dyspareunia;
- 3) is diagnosed on the basis of reported-by-the-patient fishy smell that aggravates after the intercourse;
- 4) is related to the increase in the number of Lactobacilli;
- 5) is characterized by the presence of clue cells in the smear.

The correct answer is:

- A.** 1,3.
- B.** 1,5.
- C.** 1,3,5.
- D.** 2,3,4.
- E.** 2,3,5.

Pytanie nr 51

In female adolescents without signs of breast development gonadotropins should be measured in order to exclude:

- A. hyperprolactinemia.
- B. hypergonadotropic hypogonadism.
- C. hyperthyroidism.
- D. precocious puberty.
- E. polycystic ovarian syndrome.

Pytanie nr **52**

An **absolute contraindication** to laparoscopic surgery in gynecology **is not**:

- A. patient's lack of consent.
- B. condition after extensive oncological surgery of the stomach and intestines.
- C. diffuse peritonitis.
- D. obesity.
- E. pulmonary hypertension.

Pytanie nr **53**

Ectopic pregnancy relates to the following situations:

- 1) fallopian pregnancy (ampullary region);
- 2) abdomen pregnancy;
- 3) uterine horn pregnancy;
- 4) intrauterine pregnancy;
- 5) caesarean section scar pregnancy.

The correct answer is:

- A. 1,2,3,5.
- B. 2,3,4,5.
- C. 3,4,5.
- D. 4,5.
- E. only 5.

Pytanie nr **54**

Vaginal bleeding and hypogastric pain occurred in a 25-year-old patient in 8th week of gestation. Gynecological examination showed the thinning of the cervix, and cervical dilation with parts of the gestational sac visible in the internal orifice. Indicate the correct diagnosis and the management of choice:

- A.** missed miscarriage, awaiting for the spontaneous miscarriage for 2 weeks.
- B.** inevitable miscarriage, curettage of the uterine cavity with histopathological examination.
- C.** threatened miscarriage, bed rest.
- D.** anembryonic pregnancy, awaiting for the spontaneous miscarriage for 2 weeks.
- E.** missed miscarriage, intravaginal therapy with progesterone.

Pytanie nr **55**

Indicate true statements concerning male reproductive system:

- 1) cycle of the seminiferous epithelium takes approx. 74 days;
- 2) in the testes FSH acts mainly on Sertoli and LH mainly on Leydig cells;
- 3) after gonadotropin stimulation Leydig cells produce testosterone;
- 4) estrogens do not participate in the regulation of male reproductive system.

The correct answer is:

- A.** all the above.
- B.** 2,4.
- C.** 1,2,3.
- D.** 2,3,4.
- E.** 3,4.

Pytanie nr **56**

Which of the following is called menarche?

- A.** first menstruation in life, usually occurring at the age of 12-13 years.
- B.** development of pubic hair, usually occurring at the age of 11 years.

- C. beginning of the hormonal activity of the ovaries.
- D. evolving function of the adrenals that comes 2-3 years before gonadal function.
- E. development of the breasts, usually occurring at the age of 10 years.

Pytanie nr 57

A basic recommendation for a woman with polycystic ovary syndrome, BMI>27, irregular menstrual cycles and infertility is:

- A. weight loss (diet).
- B. metformin.
- C. stimulation of ovulation with clomiphene citrate.
- D. metformin and the stimulation of ovulation with clomiphene citrate.
- E. progestogens in the second phase of the cycle.

Pytanie nr 58

Indicate the true sentences regarding screening for the neoplastic disorders of female genital organs:

- 1) screening for vaginal and vulvar cancer is not performed because of the low incidence of these tumours;
- 2) screening for endometrial cancer is not performed because its first symptoms, i.e. uterine bleeding, are easily noticed by women and reported to their doctors;
- 3) screening for non-epithelial tumours of the ovaries, the uterus and the vagina is not performed because of the low incidence of these tumours;
- 4) screening for endometrial cancer is performed and includes ultrasound imaging, the measurement of the markers: CA 125, HE 4 (human epididymis protein 4) and chorion gonadotropin;
- 5) screening for cervical cancer is performed and includes a cytological test, a test which sensitivity reaches 85-90% in leading centers.

The correct answer is:

- A. 1,2,3.
- B. 1,5.
- C. 3,4,5.

D. 4,5.

E. 1,3,4.

Pytanie nr 59

Which of the following heart diseases in a pregnant woman are absolute indications for cesarean delivery?

- 1) uncorrected ASD;
- 2) uncorrected VSD;
- 3) uncorrected coarctation of the aorta;
- 4) Marfan syndrome.

The correct answer is:

A. 2,3,4.

B. 1,2,3.

C. only 4.

D. 3,4.

E. all the above.

Pytanie nr 60

Hemorrhage, strong hypogastric pain, no palpable fundus of the uterus above the pubic symphysis and rapid deterioration of the patient's general condition during the third stage of delivery are the symptoms and signs of:

A. uterine inversion.

B. uterine rupture.

C. rupture of the vaginal fornix.

D. afterbirth retained in the uterine cavity.

E. incarcerated placenta.

Pytanie nr 61

What is the role of estrogens in the combined contraceptive pill?

- A.** fallopian tube peristalsis reduction.
- B.** thickening of cervical mucus.
- C.** inhibition of selection and growth of the dominant follicle.
- D.** suppression of the LH secretion.
- E.** A and C are correct.

Pytanie nr **62**

A 34-year-old female patient 3 months after a physiological childbirth suddenly suffered a pain of the vulva. She complained that she could not sit, and while she tries to sit down she feels a resistance in the right side. In the light of reported symptoms one should:

- A.** diagnose complications of crotch sewing and refer the patient for crotch plastic surgery.
- B.** diagnose viral vulvitis, most probably herpetic and order acyclovir therapy.
- C.** diagnose vulva carcinoma and order biopsy.
- D.** diagnose Skene's gland abscess and order antibiotics.
- E.** diagnose Bartholin's abscess, make incision and drainage.

Pytanie nr **63**

Which of the following are used to assess the maturity of girls during puberty?

- 1) Bishop score;
- 2) Nugent score;
- 3) POP-Q score;
- 4) Tanner stages;
- 5) TBS (the Bethesda system) classification;
- 6) Lauritzen criteria.

The correct answer is:

- A.** 1,4.
- B.** 2,5.
- C.** 3,5.

D. 4,5.

E. 4,6.

Pytanie nr 64

In the ultrasound examination of the fetus performed between the 11th and 13th week of pregnancy, the most accurate method of assessing the duration of pregnancy is the measurement of:

A. biparietal diameter (BPD).

B. femur length (FL).

C. abdominal circumference (AC).

D. crown-rump length (CRL).

E. gestational sac diameter.

Pytanie nr 65

Side-effects of the treatment of premenstrual syndrome with selective serotonin reuptake inhibitors include:

A. hirsutism.

B. loss of hair.

C. abnormal bleeding from the uterus.

D. inhibition of ovulation.

E. decreased libido.

Pytanie nr 66

Azoospermia is defined as:

A. absence of semen.

B. absence of sperm in semen.

C. absence of sperm mobility.

D. sperm with abnormal morphology.

E. number of sperm in semen lower than 1 million/ml.

Pytanie nr 67

Ultrasonographic examination performed at 6th week of gestation revealed coexistence of two fetuses and a single yolk sac. That allowed for the diagnosis of:

A. monochorionic, diamniotic twin pregnancy.

B. monochorionic, monoamniotic twin pregnancy.

C. dichorionic twin pregnancy, conjoint twins.

D. dichorionic twin pregnancy, with abnormal development of one of the fetuses.

E. dichorionic, monoamniotic twin pregnancy.

Pytanie nr 68

Which of the following are among factors increasing the risk of hyperemesis gravidarum?

1) cigarette smoking;

2) bulimia;

3) obesity;

4) pregnancy in a nullipara;

5) lipid metabolism disorders.

The correct answer is:

A. 1,3,5.

B. 2,3,5.

C. 2,3,4,5.

D. 1,2,3,4.

E. all the above.

Pytanie nr 69

A woman with severe preeclampsia suddenly develops respiratory depression after the administration of hydralazine and magnesium sulfate. Respiratory depression is due to:

- A.** impending eclampsia.
- B.** hydralazine toxicity.
- C.** magnesium sulfate toxicity.
- D.** pulmonary embolism.
- E.** heart attack.

Pytanie nr **70**

External examination of a woman giving birth is performed with the use of:

- 1) first Leopold's maneuver;
- 2) pelvimeter;
- 3) fifth Leopold's maneuver;
- 4) McRoberts maneuver;
- 5) Bracht maneuver.

The correct answer is:

- A.** 1,2,3.
- B.** 4,5.
- C.** 1,2.
- D.** 1,2,4.
- E.** all the above.

Pytanie nr **71**

A 42-year-old patient came to the gynecologist to get contraception advice. She smokes 5 cigarettes daily and is treated for arterial hypertension. Periodic bleeding are abundant and last up to 7 days. Which of the following methods should be advised?

- A.** progestogen-only 'mini pill'.
- B.** subcutaneous implant releasing etonogestrel.
- C.** intrauterine device with copper, Cu-IUD.
- D.** intrauterine device releasing levonorgestrel.

E. sterilization.

Pytanie nr 72

Which of the following was the most frequently occurring malignant tumor in Polish women for the 5 years (2010-2014)?

A. breast cancer.

B. endometrial cancer.

C. cervical cancer.

D. ovarian cancer.

E. lung cancer.

Pytanie nr 73

Indicate findings present in normal cardiotocographic non-stress test (NST) in pregnancy:

- 1) baseline fetal heart rate - 100/min;
- 2) baseline fetal heart rate - 150/min;
- 3) marked variability of the fetal heart rate;
- 4) absence of accelerations;
- 5) absence of decelerations.

The correct answer is:

A. 1,3.

B. 2,4.

C. 2,5.

D. 2,3.

E. 3,5.

Pytanie nr 74

A 28-year-old woman, with menstrual cycle length of about 35 days, comes to the admission room with pains in the hypogastrium and vaginal bleeding lasting for 3 days. The last menstrual flow occurred 45 days before. The patient also complains of frequent urination and tenderness of the breasts. What diagnosis should be considered first?

- A. hemorrhagic ovarian cyst.
- B. ovarian torsion.
- C. extrauterine pregnancy.
- D. adnexitis.
- E. ruptured ovarian cyst.

Pytanie nr 75

The **absolute contraindications** to the use of two-component hormonal contraceptives include:

- 1) smoking tobacco at the age over 35 years;
- 2) pregnancy or not excluded possibility of pregnancy;
- 3) migraine headaches;
- 4) mitral valve prolapse;
- 5) estrogen-dependent neoplasms.

The correct answer is:

- A. 1,2,3,4.
- B. 3,4,5.
- C. 1,2,3.
- D. 1,2,5.
- E. all the above.

Pytanie nr 76

Indicate the true sentences regarding impending miscarriages:

- 1) painless vaginal bleeding is usually present;
- 2) gynecological examination shows cervical ripening;
- 3) gynecological examination shows the cervix dilated to 1-2 cm;
- 4) speculum examination shows the amniotic sac;
- 5) treatment of choice is conservative.

The correct answer is:

- A. 1,2.
- B. 3,4.

C. 2,5.

D. 1,2,5.

E. 1,3,5.

Pytanie nr 77

As a consequence of physiological processes in the circulatory system of a pregnant woman in the case of single pregnancy the volume of circulating blood can increase by up to:

A. 20%.

B. 30%.

C. 40%.

D. 50%.

E. in a single pregnancy there is no change in the volume of circulating blood of the pregnant woman.

Pytanie nr 78

Which of the following is a risk factor for ovarian hyperstimulation syndrome?

A. long time of infertility treatment.

B. young age.

C. hypothyroidism.

D. obesity.

E. supplementation of progesterone in the luteal phase.

Pytanie nr 79

Which of the following **is not** the symptom related to uterine fibroids?

A. menorrhagia.

B. prolonged menstruation.

C. hypogastric pains.

D. pressure on the bladder or the rectum.

E. secondary amenorrhea.

Pytanie nr **80**

Indicate the true statements concerning oxytocin test:

1) it is a stress test;

2) it is a biophysical test assessing the placental efficiency;

3) positive result is concluded on the basis of late decelerations present after at least half of the contractions;

4) it is the test of choice in pregnant woman with placenta praevia.

The correct answer is:

A. all the above.

B. 1,3,4.

C. only 4.

D. 1,2,3.

E. only 1.

Pytanie nr **81**

Indicate **the false** statement concerning HPV infection:

A. so-called oncogenic HPV may cause cancers of the anus, the perineum, the penis and the oral cavity.

B. use of condoms fully eliminates the risk of HPV infection.

C. cytological result of LSIL suggests HPV infection.

D. vaccination against HPV is recommended for young girls who have not started sexual activity.

E. HPV infection does not lead to stable acquired immunity against subsequent infections with HP viruses.

Pytanie nr **82**

Indicate the true statement concerning the risk factors for ovarian cancer:

- A.** *BRCA2* mutation is more common than *BRCA1* mutation in Poland.
- B.** risk of the disease is higher in the bearers of *BRCA1* mutation than in the bearers of *BRCA2* mutation.
- C.** risk of the disease is similar in the bearers of *BRCA1* mutation and the bearers of *BRCA2* mutation.
- D.** family history of breast cancer does not increase the risk of the disease.
- E.** long-term use of hormonal contraceptives increases the risk of the disease.

Pytanie nr **83**

Routine seminological examination is an essential element in assessing the male factor as the cause of infertility. The values of microscopic evaluation of semen are presented below:

- 1) volume from 1.5-6 mL;
- 2) sperm concentration > 15 million/mL;
- 3) viability > 38% live sperm;
- 4) mobility A and B > 32%;
- 5) > 4% sperm with normal structure.

Indicate the parameters within normal range (normospermia) according to the WHO 2010 standards:

- A.** 1,2,4,5.
- B.** 2,3,4,5.
- C.** 4,5.
- D.** only 5.
- E.** 1,5.

Pytanie nr **84**

In patients with polycystic ovary syndrome there is an increased risk of:

- A.** type 2 diabetes mellitus.

- B. hypertension.
- C. endometrial cancer.
- D. myocardial infarction.
- E. all the above.

Pytanie nr 85

Indicate the true description of the influence of chorionic gonadotropin on the thyroid function:

- A. there is no influence.
- B. it lowers the TSH production which causes the decrease in thyroxine synthesis in the thyroid gland.
- C. it stimulates TSH production which causes the increase in thyroxine synthesis in the thyroid gland.
- D. it decreases TSH production and stimulates the thyroid gland to produce thyroxine.
- E. it increases TSH production and stimulates the thyroid gland to produce thyroxine.

Pytanie nr 86

A **contraindication** to sexual activity in pregnancy **is not**:

- A. multiple pregnancy.
- B. cervical incompetence.
- C. placenta previa.
- D. inflammatory urinary tract.
- E. fetal breech presentation.

Pytanie nr 87

A 25-year-old woman in 12th week of her first pregnancy made a test of antibodies against *Toxoplasma gondii*. The results were as follows: IgG = 300 IU/mL, IgM (-). A similar test performed 2 years earlier showed: IgG = 480 IU/mL, IgM (-). The patient should be

informed that:

- A.** there is a high risk of an infection with *Toxoplasma gondii* caused by the consumption of both raw meat and contaminated fruit and vegetables.
- B.** she has to repeat the test of antibodies against *T. gondii* every 6-8 weeks during the whole pregnancy.
- C.** she should undergo antiprotozoal treatment.
- D.** she was infected with *Toxoplasma gondii* before the pregnancy (most probably it was a subclinical infection) and her child is not at risk of the infection.
- E.** it is necessary to perform invasive diagnostics for the infection (amniocentesis, examination of the amniotic fluid for protozoan cysts).

Pytanie nr **88**

Which of the following statements concerning hydatidiform mole is **false**?

- A.** complete hydatidiform mole is diploid and all the genetic material is of paternal origin.
- B.** incomplete hydatidiform mole is usually triploid.
- C.** principal diagnostic tool for hydatidiform mole is ultrasound imaging.
- D.** first symptom of hydatidiform mole is usually uterine bleeding.
- E.** most common symptoms of hydatidiform mole are excessive vomiting and the symptoms of hypothyroidism.

Pytanie nr **89**

Oligohydramnios is defined as the decrease in the volume of amniotic fluid below:

- A.** 600 mL.
- B.** 500 mL.
- C.** 400 mL.
- D.** 300 mL.
- E.** 200 mL.

Pytanie nr 90

Which sexually transmitted disease **does not** cause symptoms in the genital area?

- A. infection caused by HIV.
- B. infection caused by HPV.
- C. infection caused by HSV.
- D. syphilis.
- E. gonorrhoea.

Pytanie nr 91

A 41-year-old patient came to gynecologist asking for advice on contraception during perimenopause. She has two children, and has diabetes mellitus type I diagnosed at the age of 4. The current level of glycated haemoglobin HbA_{1c} is 8.9%, urine albumin excretion - 40 mcg/mg creatinine, her blood pressure is about 130/80 mmHg, ophthalmoscopy showed pre-proliferative retinopathy. Which of the following methods is **contraindicated** in this patient?

- A. cervical mucus-body temperature method.
- B. progestin-only pill.
- C. combined oral contraceptive pills with ethinylestradiol and levonorgestrel.
- D. barrier method.
- E. intrauterine device with levonorgestrel.

Pytanie nr 92

Which of the following is the correct management of a 30-year-old patient in 37th week of her first pregnancy who was diagnosed with the minor type of placenta previa confirmed by repeated US examination and reports minor bleeding from the reproductive tract?

- A. transvaginal palpation, transvaginal ultrasound examination, conservative treatment (pregnancy continuation up to 38 weeks).
- B. transvaginal palpation, CTG monitoring, vaginal delivery.

C. assessment of the general condition of the mother, fetal pulse monitoring, preparations and performing caesarean section.

D. assessment of the general condition of the mother, fetal pulse monitoring, vaginal delivery.

E. speculum examination, intravenous fenoterol infusion, pregnancy continuation up to 38 weeks.

Pytanie nr 93

Prohibited vaccinations during pregnancy include:

A. measles.

B. hepatitis B.

C. influenza.

D. tetanus.

E. yellow fever.

Pytanie nr 94

A 28-year-old woman who fainted again at home reported to the obstetrics and gynecology ER. On admission she presented pale, sweaty face, rapid heart rate, body temperature 36.6°C. Despite inserting a contraceptive intrauterine device a year ago, she had her last period 8 weeks ago. Preliminary gynecological examination showed: slight vaginal bleeding, slightly enlarged uterus, painful resistance in the projection of the left uterine appendages and doughy tumor in the pouch of Douglas. These symptoms are indicative of diagnosis:

A. ectopic pregnancy - tubular miscarriage.

B. cervical pregnancy.

C. correctly located pregnancy - incomplete miscarriage.

D. acute adnexitis.

E. ovarian cancer.

Pytanie nr 95

Choriocarcinoma is one of the rarest but most aggressive malignant tumors in women. Its diagnosis is based on:

- A.** magnetic resonance imaging of the abdominal cavity and pelvis.
- B.** histopathological result from biopsy or curettage of the uterine cavity.
- C.** histopathological result obtained during hysterectomy.
- D.** high levels of hCG in blood serum.
- E.** pregnancy induced hypertension before the 20th week of pregnancy.

Pytanie nr 96

Which of the following increases the risk of monozygotic twin pregnancy?

- A.** advanced maternal age.
- B.** use of ovulation stimulating drugs.
- C.** previous multiple pregnancy.
- D.** long-term use of oral contraceptives.
- E.** none of the above.

Pytanie nr 97

A 58-year-old patient complains of vaginal dryness and postcoital bleedings. The gynecological examination reveals atrophic lesions in the vagina and normal uterus and uterine appendages. The cytological smear of the cervix is normal. The local vaginal estrogen therapy has been recommended and thus the patient should:

- A.** receive simultaneous gestagen therapy in order to prevent the overgrowth of endometrium and the development of endometrial cancer.
- B.** receive simultaneous gestagen therapy in order to prevent the development of breast cancer.
- C.** receive simultaneous gestagen therapy in order to prevent the development of cervical cancer.

D. receive simultaneous gestagen therapy in order to decrease the risk of a stroke.

E. there is no indication for gestagen therapy in this patient.

Pytanie nr **98**

Complete androgen insensitivity syndrome is characterized by:

A. female phenotype, no axillary hair, male external genitalia.

B. male phenotype, no pubic hair, male external genitalia.

C. female phenotype, pubic and axillary hair, female external genitalia.

D. 46 XY karyotype with estrogen-secreting ovaries.

E. female phenotype, no pubic and axillary hair, female external genitalia.

Pytanie nr **99**

Which of the following **is not** an advantage of Pfannenstiel incision?

A. good cosmetic effect.

B. rare cases of wound dehiscence.

C. wide access to the operative field.

D. rare cases of incisional hernia.

E. better blood supply to the wound.

Pytanie nr **100**

Which of the following **is not** a biological effect of progesterone?

A. induction of endometrial secretory changes during the ovulation cycle.

B. cervical mucus change into impenetrable to sperm.

C. decrease of the hypoglycemic effect of insulin.

D. decrease of contractibility of the uterine muscle and fallopian tube peristalsis.

E. increase in the concentration of coagulation factors II, VII, IX and X.

Pytanie nr **101**

The composition of the amniotic fluid includes the following proteins produced by the foetus:

- 1) transferrin;
- 2) alpha-fetoprotein;
- 3) beta-microglobulin;
- 4) IgG immunoglobulins.

The correct answer is:

- A.** 1,2,4.
- B.** 1,2,3.
- C.** 2,3,4.
- D.** only 2.
- E.** all the above.

Pytanie nr **102**

Which of the following is the primary medication used for the long-term antihypertensive therapy in pregnancy?

- A.** methyldopa.
- B.** perindopril.
- C.** enalapril.
- D.** ramipril.
- E.** lisinopril.

Pytanie nr **103**

An umbilical cord prolapse during the second stage of the labor in a cephalic presentation with a longitudinal lie:

- A.** is not dangerous.
- B.** is an indication for the attempt of a manual reduction of the prolapse.

- C. is an indication for immediate delivery.
- D. is an indication for a manual rotation of the fetus.
- E. is an indication for the immediate artificial rupture of membranes.

Pytanie nr **104**

Fetal bradycardia is defined as a decrease of the basic fetal heart rate:

- A. below 100 beats/min. lasting at least 15 minutes.
- B. below 110 beats/min. lasting at least 10 minutes.
- C. below 110 beats/min. lasting at least 15 minutes.
- D. below 120 beats/min. lasting at least 10 minutes.
- E. below 120 beats/min. lasting at least 15 minutes.

Pytanie nr **105**

Which of the following factors **does not** predispose to the breech presentation?

- A. oligohydramnios.
- B. polyhydramnios.
- C. localization of the placenta on the posterior wall of the uterus.
- D. hydrocephalus.
- E. placenta praevia.

Pytanie nr **106**

The administration of short-acting GnRH analogues begins on the first day of the menstrual cycle and from the third day gonadotropins are added. This description regards controlled ovarian hyperstimulation with the use of:

- A. ICSI technique (intracytoplasmic sperm injection).
- B. slow freezing of reproductive material.

C. protocol with GnRH antagonists.

D. long protocol.

E. short protocol.

Pytanie nr **107**

Which of the statements regarding stress urinary incontinence is true?

A. increase in intra-abdominal pressure is accompanied by involuntary loss of urine.

B. involuntary leakage of urine is accompanied by a strong but short urinary urgency.

C. volume of leaking urine is large.

D. frequency of micturition at night is increased.

E. in urodynamic testing detrusor hyperreflexia is observed.

Pytanie nr **108**

Endometrial cancer is currently the most common female genital cancer in developed countries. The following are the characteristics of its two subtypes:

1) estrogen dependence (positive E + and P + receptor state);

2) aggressive course, frequent metastases;

3) the most common are mutations PTEN, K-ras, microsatellite instability;

4) the most common histopathological type is endometrioid endometrial cancer;

5) the most common mutations are p-53, HER2-neu, p16.

The correct combination of characteristics of the first subtype of endometrial cancer is:

A. 1,2,5.

B. 2,4,5.

C. 3,4.

D. 1,3,4.

E. only 1.

Pytanie nr **109**

A disulfiram-like reaction can be expected in a patient who was drinking alcohol while

undergoing the treatment for the symptoms of a genital infection with:

- A. HPV virus.
- B. HSV virus.
- C. *Chlamydia trachomatis*.
- D. *Trichomonas vaginalis*.
- E. *Neisseria gonorrhoeae*.

Pytanie nr 110

Which of the following are maternal risk factors for the breech position?

- 1) defects of the uterus which impair fetal rotation;
- 2) leiomyomas of the uterus;
- 3) narrow pelvis;
- 4) excessively wide pelvis;
- 5) multiparity.

The correct answer is:

- A. 1,3.
- B. 1,2,4.
- C. 1,3,5.
- D. all the above.
- E. only 5.

Pytanie nr 111

The complications of fetal diabetes include:

- A. macrosomia.
- B. preterm birth, respiratory distress syndrome.
- C. hypoglycemia, polycythemia, hypokalemia.
- D. all the above.

E. none of the above.

Pytanie nr 112

Uterine leiomyomas are the most frequent benign neoplasms of the uterus. Indicate **the false** statement concerning their symptoms:

- A.** they include prolonged heavy periods and intermenstrual bleedings.
- B.** they include pain in the hypogastrium, and pressure on the bladder or the rectum.
- C.** small leiomyomas may be asymptomatic.
- D.** hot flushes are typical symptoms.
- E.** leiomyomas may be the cause of infertility.

Pytanie nr 113

The diagnosis of endometriosis should be considered in a patient with the following symptoms:

- 1) periodic pain, beginning just before the menstrual period lasting throughout the whole time of menstruation;
- 2) the temperature during the menstrual period $> 39^{\circ}\text{C}$;
- 3) dyspareunia (pain related to sexual intercourse);
- 4) pain during defecation or micturition;
- 5) a stinking discharge from the vagina, occurring cyclically after menstruation.

The correct answer is:

- A.** 2,3,4.
- B.** 1,3,4.
- C.** 1,3,4,5.
- D.** 2,5.
- E.** all the above.

Pytanie nr 114

The most common site for an ectopic pregnancy is:

- A.** abdominal cavity.

- B. ovary.
- C. intramural part of the uterine tube.
- D. isthmus of the uterine tube.
- E. ampulla of the uterine tube.

Pytanie nr **115**

Intermittent amenorrhea denotes:

- A. lack of menstruation under 16 years of age.
- B. menstrual bleeding lasting 5-7 days.
- C. menstrual bleeding occurring every 43 days - 6 months.
- D. menstrual bleeding occurring with gaps shorter than 21 days.
- E. alternative name for coitus interruptus.

Pytanie nr **116**

Indicate the true sentences regarding ovarian reserve:

- 1) it is lower in Afro-Americans than in Caucasian women;
- 2) it is determined genetically;
- 3) it decreases after ovarian surgery;
- 4) it decreases as a consequence of ovarian disorders;
- 5) it increases with woman's age;
- 6) it decreases after antibiotic treatment.

The correct answer is:

- A. 1,2,3.
- B. 2,3,4.
- C. 3,4,5.
- D. 4,5,6.
- E. 1,4,5.

Pytanie nr 117

In a 16-year-old girl with amenorrhea, vaginal hematoma (hematocolpos) was diagnosed during ultrasound examination. The most probable diagnosis is:

- A. imperforate hymen.
- B. uterine septum.
- C. uterine aplasia.
- D. Mayer-Rokitansky-Küster-Hauser syndrome.
- E. ovarian agenesis.

Pytanie nr 118

Absence of menstruation after fertilization and implantation of the embryo in the endometrium is caused by:

- A. absence of corpus luteum degeneration and the increased synthesis of progesterone and estrogens because of increasing levels of chorionic gonadotropin.
- B. absence of corpus luteum degeneration and the increased synthesis of progesterone and estrogens because of decreasing levels of prolactin.
- C. absence of corpus luteum degeneration because of the placenta taking over the production of progesterone and estrogens.
- D. degeneration of the corpus luteum because of the placenta taking over the production of progesterone and estrogens.
- E. degeneration of the corpus luteum because of increasing prolactin levels.

Pytanie nr 119

Which of the following medicines **is not** indicated in endometriosis treatment?

- A. danazol.
- B. goserelin.
- C. mirabegron.

D. dienogest.

E. progesteron.

Pytanie nr 120

The basic strategy of treatment in the case of Bartholin's gland abscess consists in:

A. incision of the abscess or its marsupialization.

B. surgical removal of the abscess.

C. observation.

D. antibiotic therapy only.

E. wraps with washing soda.

Pytanie nr 121

Is it acceptable to postpone the next dose when taking contraceptive tablets containing only progestogen (mini pills)?

A. yes, up to 3 hours.

B. yes, up to 12 hours.

C. yes, up to 24 hours.

D. yes, up to 72 hours.

E. the next dose should absolutely be taken at the same time.

Pytanie nr 122

Oral hormonal contraception is a well-documented protective factor against:

1) ovarian cancer;

2) liver cancer;

3) endometrial cancer;

4) cervical cancer;

5) renal cancer.

The correct answer is:

A. 2,3,4.

B. 1,5.

C. 1,3,4.

D. 1,3.

E. 2,4,5.

Pytanie nr 123

Pelvic floor disorders are the result of the weakening of the supporting and hanging apparatus as well as an excessive action of forces causing a displacement of genital organs. The factors conducive to the decline of the reproductive organ include:

- 1) damage to the vulvar and pelvic nerves in the course of vaginal delivery;
- 2) physical work in standing position and associated with carrying;
- 3) respiratory diseases associated with chronic cough;
- 4) chronic constipation;
- 5) post-menopausal estrogen deficiency.

The correct answer is:

A. 1,2,3,4.

B. 1,3,4,5.

C. 1,2.

D. only 1.

E. all the above.

Pytanie nr 124

A 25-year-old woman was referred to the GP because of secondary amenorrhea. She reports galactorrhoea and periodically blurred vision. Which of the following hormones should be measured?

A. adrenocorticotropic.

B. glucagon.

C. insulin.

D. prolactin.

E. testosterone.

Pytanie nr 125

Indicate the true statements concerning the double test in prenatal diagnostics:

- 1) it is performed between 11th and 14th week of gestation;
- 2) it assesses 2 biochemical markers in the blood of the pregnant woman;
- 3) in the case of a fetus with Down syndrome the concentration of free beta hCG subunit is decreased;
- 4) in the case of a fetus with Down syndrome the concentration of PAPP-A is decreased;
- 5) in the case of a fetus with Edwards syndrome the concentration of free beta hCG subunit is decreased.

The correct answer is:

- A. 1,2,3,4.**
- B. 1,3,4,5.**
- C. 2,3,4,5.**
- D. 1,2,4,5.**
- E. all the above.**

Pytanie nr 126

The left uterine artery is a direct branch of:

- A. aorta.**
- B. left common iliac artery.**
- C. left external iliac artery.**
- D. left internal iliac artery.**
- E. left ovarian artery.**

Pytanie nr 127

The most common uterine anomaly is:

- A. unicorn uterus with a rudimentary horn.**

B. uterus didelphys.

C. septate uterus.

D. bicornuate uterus.

E. unicorn uterus.

Pytanie nr 128

Labor begins with regular contractions of the uterus causing the cervix to shorten and open. Which of the following concerning the first stage of labor is true?

- 1) it is the time from full cervical dilation to the birth of a child;
- 2) it is the time of postpartum content expulsion;
- 3) it is the time from the beginning of labor to the full cervical dilation;
- 4) it is divided into two phases: latent - from the labor beginning to the opening around 3-4 cm and active - from the 3-4 cm of the dilation to the complete dilation;
- 5) maximal time of the first stage of labor is 2 hours.

The correct answer is:

A. only 1.

B. only 2.

C. 3,4.

D. 3,4,5.

E. none of the above.

Pytanie nr 129

“This tumor can develop before puberty but the peak incidence is observed in women of reproductive age. It easily ruptures which may lead to peritoneal signs and symptoms demanding urgent surgery. In postmenopausal women it may cause abnormal bleeding from the reproductive tract. Surgical excision does not guarantee full recovery as this tumor recurrence may be observed as many as 30 years after the initial diagnosis”. The above is a characteristic of:

A. ovarian cancer.

B. sarcoma.

- C. hydatidiform mole.
- D. germinoma.
- E. granulosa cell tumor.

Pytanie nr **130**

Azoospermia is defined as:

- A. absence of spermatozoa in semen.
- B. absence of semen.
- C. absence of progressive motility of sperms.
- D. concentration of sperms less than 20 million in 1 mL of semen.
- E. less than 30 % of spermatozoa with normal morphology.

Pytanie nr **131**

Which of the following **is not** a risk factor for urinary incontinence?

- A. patient's age.
- B. obesity.
- C. previous gynecological procedures.
- D. nulliparity.
- E. chronic constipation.

Pytanie nr **132**

The combination of pharmacological therapy with a planned cesarean delivery has been shown to reduce the perinatal transmission of HIV to:

- A. 25%.
- B. 17.5%.
- C. 5%.

D. 1.5%.

E. perinatal transmission of HIV does not depend on the mode of a delivery.

Pytanie nr 133

The main source of alfa-fetoprotein (AFP) in the maternal blood at 16 weeks' gestation is:

A. maternal liver.

B. fetal liver.

C. fetal adrenal glands.

D. placenta.

E. yolk sac.

Pytanie nr 134

Pregnancy in a woman with Swyer's syndrome may be possible:

A. after stimulation of ovulation with gonadotropins.

B. after performing patient's oocyte retrieval by the aspiration of ovarian follicles.

C. when a donor oocyte is used.

D. after surgical recreation of the vagina.

E. after the uterus transplantation.

Pytanie nr 135

Indicate the **false** statement on interpretation of CTG:

A. sporadic accelerations are indicative of a healthy foetus.

B. bradycardia within 100-110 bpm does not indicate that foetal health is endangered.

C. cardiotocography should be implemented when the foetus is mature enough to survive outside the womb.

D. decelerations are drops in the baseline heart rate lasting 20 seconds or longer with a magnitude of at least 10 bpm.

E. early decelerations mirror uterine contractions.

Pytanie nr **136**

Indicate the estimated date of delivery calculated according to Naegele's rule for a woman with regular menstrual cycles of 32 days if the last menstruation started on 15 July 2018:

A. 15 April 2019.

B. 19 April 2019.

C. 21 April 2019.

D. 25 April 2019.

E. 26 April 2019.

Pytanie nr **137**

A patient presents at the gynecological outpatient clinic on the 12th day after the radical hysterectomy due to endometrial hyperplasia with atypia. She complains of a constant outflow of urine from the vagina. Which of the following should be suspected?

A. overactive bladder.

B. exercise-induced urinary incontinence caused by the impairment of ligaments of the uterus.

C. mixed urinary incontinence.

D. evacuation of accumulated lymph fluid through the vaginal stump.

E. vesicovaginal fistula.

Pytanie nr **138**

Factors increasing the risk of emergency peripartum hysterectomy **do not** include:

A. previous cesarean section.

B. pregnancy after in vitro procedure.

C. placenta praevia or percreta.

D. numerous previous pregnancies.

E. age above 35.

Pytanie nr **139**

The surgical treatment of cervical incompetence consists in a cervical cerclage. Which of the following conditions should be met before putting on a cerclage?

- 1) normal fetal development;
- 2) alive fetus;
- 3) absence of uterine contractions;
- 4) normal vaginal biocenosis;
- 5) gestational age over 28 weeks.

The correct answer is:

A. 2,3,4,5.

B. 1,3,4,5.

C. 1,2,3,4.

D. all the above.

E. 1,2.

Pytanie nr **140**

A normal fetal heart rate is in the range of:

A. 160-180 beats/minute.

B. 110-160 beats/minute.

C. 100-120 beats/minute.

D. 80-100 beats/minute.

E. 60-80 beats/minute.

Pytanie nr **141**

Which of the following **is not** characteristic of thrombotic thrombocytopenic purpura in a pregnant woman?

- A. microangiopathic hemolytic anemia.
- B. disseminated intravascular coagulation.
- C. thrombocytopenia.
- D. kidney failure.
- E. fever.

Pytanie nr **142**

Obstetrical complications that may lead to disseminated intravascular coagulation (DIC) **do not** include:

- A. amniotic fluid embolism.
- B. premature placental abruption of the normally located placenta.
- C. HELLP syndrome.
- D. eclampsia.
- E. fetal hypotrophy caused by placental insufficiency.

Pytanie nr **143**

Which of the following **is not** a risk factor for endometrial cancer?

- A. BMI > 32 kg/m².
- B. tamoxifen therapy.
- C. late menopause.
- D. nulliparity.
- E. family history of ovarian cancer.

Pytanie nr **144**

The best method to diagnose fetal hemolytic disease is:

- A. amniocentesis.

- B.** cordocentesis.
- C.** fetoscopy.
- D.** placenta biopsy.
- E.** biopsy of the fetal liver.

Pytanie nr **145**

The phase of voiding the urinary bladder, initiated by the brain cortex is characterized by the relaxation of sphincter muscles surrounding the urethra and the increased tone of the detrusor muscle. These phenomena are caused by:

- A.** inhibition of the sympathetic activity and the increased pulsation in the parasympathetic system.
- B.** inhibition of the parasympathetic activity and the increased pulsation in the sympathetic system.
- C.** increased pulsation in the somatic system innervating muscles of the pelvic floor.
- D.** inhibition of the sympathetic activity only.
- E.** inhibition of the parasympathetic activity only.

Pytanie nr **146**

Fetal heart auscultation is an important part of obstetrical examination and it helps safe delivery management. During normal delivery the basic fetal heart rate should be in the range of:

- A.** 70 - 90 bits per minute.
- B.** 100 - 170 bits per minute.
- C.** 110 - 150 bits per minute.
- D.** 110 - 190 bits per minute.
- E.** above 170 bits per minute.

Pytanie nr **147**

Which of the following indicates a complete mole at ultrasound examination?

- 1) "snow-storm" image;
- 2) no fetus;
- 3) presence of theca lutein cysts;
- 4) presence of fetal echo with a visible heart action;
- 5) presence of the gestational sac with amniotic fluid.

The correct answer is:

- A.** 1,2,3.
- B.** 1,2,5.
- C.** 1,3,5.
- D.** 2,3,5.
- E.** 3,4,5.

Pytanie nr 148

A 65-year-old patient in good medical condition, with arterial hypertension and type 2 diabetes mellitus treated with oral therapy, comes to the gynecologist because of the moderate hemorrhage from the reproductive tract. Her body weight is 87 kg and the height is 163 cm. The result of the endometrial biopsy is: endometrial adenocarcinoma G1. Transvaginal ultrasound imaging suggests minimal invasion of the endometrium, CT imaging shows no intra-abdominal dissemination nor pathological pelvic or paraaortic lymph nodes. Indicate the optimal treatment for this patient:

- A.** surgery: hysteroscopic removal of the tumor.
- B.** surgery: panhysterectomy.
- C.** radiotherapy (brachytherapy + teletherapy).
- D.** chemotherapy (paclitaxel + carboplatin).
- E.** hormonal (megestrol acetate).

Pytanie nr 149

Indicate the **false** sentence concerning the influence of progesterone on the female body:

- A.** progesterone has a diuretic effect.

- B.** progesterone thickens cervical mucus.
- C.** progesterone increases glycogen synthesis.
- D.** progesterone increases the thickness of the myometrium and the rate of peristaltic waves of the fallopian tubes.
- E.** progesterone increases body temperature.

Pytanie nr **150**

Indicate the true statement concerning vulvar cancer:

- A.** most often occurs before the age of 40.
- B.** accounts for 3-8% of all female genital malignancies.
- C.** more than 90% of vulvar cancers are glandular cancers.
- D.** the treatment of choice is a systemic treatment.
- E.** five-year survival for women in FIGO stage I is 31%.

Pytanie nr **151**

Which of the following manoeuvres **is not** recommended in the case of shoulder dystocia:

- A.** suprapubic pressure.
- B.** Zavanelli's manoeuvre.
- C.** Bracht's manoeuvre.
- D.** McRoberts manoeuvre.
- E.** Woods manoeuvre.

Pytanie nr **152**

Indicate the correct sequence of fetal head movements in the birth canal during labor:

- A.** flexion, extension, rotation, rotation.
- B.** extension, rotation, flexion, rotation.

C. flexion, rotation, rotation, extension.

D. rotation, flexion, extension, rotation.

E. flexion, rotation, extension, rotation.

Pytanie nr **153**

In which of the following neoplasms brachytherapy **is not** used as an auxiliary method?

A. ovarian cancer.

B. endometrial cancer.

C. cervical cancer.

D. vulvar cancer.

E. in the all of the above cancers brachytherapy is used as an auxiliary method.

Pytanie nr **154**

Finding the lambda sign on ultrasound examination in a patient at 9 week of gestation allows diagnosing:

A. missed miscarriage.

B. monoamniotic twin pregnancy.

C. monozygotic twin pregnancy.

D. singleton pregnancy.

E. dichorionic twin pregnancy.

Pytanie nr **155**

Which of the following **does not** affect the result of the triple test?

A. body weight of the pregnant woman.

B. tobacco smoking.

C. number of pregnancies.

D. race.

E. type I diabetes.

Pytanie nr **156**

Which of the following are risk factors for exercise-induced urinary incontinence?

- 1) young age;
- 2) BMI < 30 kg/m²;
- 3) chronic constipation;
- 4) taking diuretics or anti-hypertensive medications;
- 5) chronic diseases with coughing.

The correct answer is:

A. 1,2.

B. 3,4,5.

C. 1,3,5.

D. 2,4,5.

E. all the above.

Pytanie nr **157**

Routine obstetric ultrasounds between 11 weeks and 14 weeks of gestational age **do not** include:

- A. evaluation of the number of gestational sacs in the uterine cavity.
- B. measurement of the crown-rump length in the sagittal plain.
- C. evaluation of the blood flow in the umbilical artery.
- D. evaluation of the fetal heart rate (FHR).
- E. determination of the number of chorionic sacs for multiple gestations.

Pytanie nr **158**

Which of the following is the most probable diagnosis in the case of impaired fertility, chronic pelvic pain, painful periods, and dyspareunia?

- A.** endometriosis.
- B.** ovarian cancer.
- C.** cervical cancer.
- D.** adnexitis.
- E.** premature ovarian failure.

Pytanie nr **159**

Assessment of the location of the fundal height is an important part of obstetrical examination. The fundus of the uterus at 24th week of gestation is normally located:

- A.** 3 finger widths above the pubic symphysis.
- B.** at the umbiculus.
- C.** 3 finger widths above the umbiculus.
- D.** 3 finger widths below the xiphoid process.
- E.** under the costal margin.

Pytanie nr **160**

A patient at 26th week of the gestation was injured in a traffic accident and subsequently was treated surgically and received 2 medications of the category B in the FDA classification. The category B denotes:

- A.** medications with the documented risk of fetal abnormalities.
- B.** medications with the documented risk of fetal abnormalities, but benefits from their use justify their application.
- C.** medications that showed adverse effect on the fetus in animal reproduction studies and there are no adequate studies in humans, but potential benefits may warrant the use of the drug in pregnant women despite a potential risk.
- D.** medications that are relatively safe in pregnancy, with no teratogenic effects observed in animal studies, but there are no adequate studies in humans.
- E.** medications that did not show teratogenic effects in the studies in pregnant women.

Pytanie nr **161**

Which of the following **is not** a germ cell tumor of the ovary?

- A. germinoma.
- B. teratoma.
- C. embryonal carcinoma.
- D. granulosa cell tumor.
- E. choriocarcinoma.

Pytanie nr **162**

Which of the following are indications for first-trimester complex prenatal diagnosis?

- 1) advanced maternal age (at least 35 years at the expected time of delivery);
- 2) giving birth to a child with genetic disorders in the past;
- 3) increased nuchal translucency (NT) values;
- 4) preeclampsia in the previous pregnancy;
- 5) oligohydramnios.

The correct answer is:

- A. 1,4,5.
- B. 2,4,5.
- C. 1,2,3.
- D. 4,5.
- E. 3,4.

Pytanie nr **163**

A single-handed maneuver performed during palpation of the pregnant woman's abdomen that allows to determine the leading part of the fetus and its location in relation to the inlet is called:

- A. first Leopold's maneuver.
- B. second Leopold's maneuver.

- C. third Leopold's maneuver.
- D. forth Leopold's maneuver.
- E. none of the above.

Pytanie nr **164**

Fetal lie is:

- A. relation of the particular parts of the fetus to each other.
- B. relation of the particular parts of the fetus to the particular parts of the uterus.
- C. relation of the long fetal axis to the long axis of the uterus.
- D. relation of the particular parts of the fetus to the long axis of the uterus.
- E. relation of the long fetal axis to the particular parts of the uterus.

Pytanie nr **165**

Indicate the true statements concerning vulvar cancer:

- 1) most commonly it develops from the stratified squamous epithelium;
- 2) HPV infections play an important role in pathogenesis of some vulvar cancers;
- 3) supraclavicular lymph nodes are the most common site of metastasis;
- 4) basic treatment consists in the surgery and radiochemotherapy;
- 5) hormonal therapy is the most effective method of vulvar cancer treatment.

The correct answer is:

- A. only 5.
- B. 1,2,3.
- C. 1,2,5.
- D. 2,3,4.
- E. 1,2,4.

Pytanie nr **166**

A 32-year-old patient did the sperm count test for the diagnostics of couple infertility. He

avoided ejaculation for 3 days before the test. The test results show: sperm number 18 mln/mL; sperm motility slow and rapid 33% in total; sperm morphology - 5% normally shaped. What conclusion should be formulated?

- A. normospermia.
- B. asthenozoospermia.
- C. kryptozoospermia.
- D. oligoteratozoospermia.
- E. asthenoteratozoospermia.

Pytanie nr **167**

In patients with severe preeclampsia the treatment of choice, both for the cessation of seizures as well as for their prevention, includes:

- A. atosiban administered intravenously.
- B. atosiban administered intramuscularly.
- C. magnesium sulfate administered intravenously.
- D. magnesium sulfate administered intramuscularly.
- E. calcium gluconate administered intravenously.

Pytanie nr **168**

Chronic alcohol abuse may lead to disorders of spermatogenesis as the result of:

- A. increased concentration of levonorgestrel.
- B. decreased adrenal production of androstendione.
- C. increased concentrations of estrogens.
- D. increased synthesis of LH.
- E. increased synthesis of testosterone.

Pytanie nr **169**

Which of the following is characteristic of stress urinary incontinence (SUI) in women?

- A.** involuntary voiding that is preceded by strong urge to void and nycturia.
- B.** leakage of urine following increased abdominal pressure.
- C.** involuntary leakage of urine as the bladder is filled over.
- D.** continuous urinary leakage through the fistula.
- E.** feeling of incomplete voiding after emptying the bladder.

Pytanie nr 170

The secondary prophylaxis of cervical cancer consists in:

- A.** pro-healthy style of life to avoid HPV infection.
- B.** cytological screening.
- C.** cervical conization in the case of CIN2+ cytological outcome.
- D.** radical hysterectomy with lymphadenectomy in the case of invasive cancer.
- E.** brachytherapy in the case of metastatic cancer.

Pytanie nr 171

The true conjugate is defined as:

- A.** line connecting the lower edge of the pubic symphysis with the iliococcygeus joint.
- B.** line connecting the promontory with the upper edge of the pubic symphysis.
- C.** line connecting the promontory with the most inwardly facing part of the pubic symphysis.
- D.** line connecting a cavity between the spinous processes of L3 and L4 with the upper edge of the pubic symphysis.
- E.** line connecting the promontory with the lower edge of the pubic symphysis.

Pytanie nr 172

The importance of GnRH analogues in gynecology has long been known. They are used in

the following situations, **except for**:

- A.** endometriosis treatment.
- B.** in vitro fertilization programs.
- C.** uterine myoma treatment.
- D.** precocious puberty treatment.
- E.** growth hormone deficiency treatment.

Pytanie nr **173**

The clinical features of HELLP syndrome include:

- 1) elevated liver enzymes;
- 2) low platelet count;
- 3) hemolysis;
- 4) thrombocytosis;
- 5) decreased concentration of indirect bilirubin.

The correct answer is:

- A.** 1,2,3.
- B.** 1,3,4.
- C.** 1,3,5.
- D.** 2,3,5.
- E.** 3,4,5.

Pytanie nr **174**

Which of the following may cause retrograde ejaculation?

- 1) diabetes mellitus;
- 2) depression;
- 3) sclerosis multiplex;
- 4) spinal cord injury;
- 5) myocardial infarction;
- 6) femoral bone fracture.

The correct answer is:

- A.** 1,3,4.

B. 1,4,5.

C. 2,5,6.

D. 3,4,5.

E. 4,5,6.

Pytanie nr 175

The ultrasound risk factor for shoulder dystocia **is not**:

A. estimated fetal weight above 4500 g.

B. abdominal circumference above 350 mm.

C. difference in the circumference of the abdomen and fetal head over 40 mm.

D. difference in the transverse abdominal and biparietal diameter over 26 mm.

E. difference in the length of the clavicular bone and the humerus over 50 mm.

Pytanie nr 176

A heterotopic pregnancy may be a complication of an assisted reproductive program and denotes:

A. ectopic pregnancy localized in the abdominal cavity.

B. ectopic pregnancy localized in the scar after the cesarean section.

C. ectopic pregnancy localized in the isthmus of the uterine tube.

D. ectopic pregnancy localized in the cervix.

E. both extra-uterine and intrauterine pregnancy occurring simultaneously.

Pytanie nr 177

The lower limit of the reference range of hemoglobin concentration below which anemia is diagnosed in a pregnant woman is:

A. higher than in non-pregnant women because the increase in the volume of circulating blood in pregnancy does not keep up with the increase in erythrocyte production.

- B.** lower than in non-pregnant women because the increase in erythrocyte production does not keep up with the increase in the volume of circulating blood.
- C.** the same as in non-pregnant women.
- D.** irrelevant because anemia in pregnancy is diagnosed on the basis of hematocrit.
- E.** dependent on the level of hemoglobin concentration before the pregnancy.

Pytanie nr **178**

Sheehan's syndrome - postpartum pituitary gland necrosis - is characterized by:

- A.** hypogonadotropic hypergonadism.
- B.** hypogonadotropic hypogonadism.
- C.** hypergonadotropic hypergonadism.
- D.** hypergonadotropic hypogonadism.
- E.** primary amenorrhoea.

Pytanie nr **179**

Which of the following is introduced into the peritoneal cavity in order to produce pneumoperitoneum during the laparoscopic procedure?

- A.** saline solution.
- B.** carbon dioxide.
- C.** oxygen.
- D.** nitrogen.
- E.** compressed air.

Pytanie nr **180**

Which of the following markers should be used in the diagnosis of molar pregnancy?

- A.** Ca125.

B. beta HCG.

C. estradiol.

D. inhibin A.

E. HE4.

Pytanie nr **181**

Indicate the true sentence describing endometriosis:

A. retrograde menstrual flow is necessary in the pathogenesis of endometriosis.

B. surgical treatment of endometriosis should not be used in women with pelvic pain.

C. fertility in endometriosis is only reduced when adhesions in the peritoneal cavity are present.

D. pharmacological treatment in endometriosis is limited due to the side effects of medicines.

E. every woman with endometriosis is infertile.

Pytanie nr **182**

The multiple pregnancy is related to the increased risk of numerous obstetrical complications compared with the singleton pregnancy. Which of the following complications **are not** observed in multiple pregnancies more often than in singleton pregnancies?

1) shoulder dystocia;

2) gestational hypertension;

3) necessity for the cesarean section;

4) postterm pregnancy;

5) perinatal death;

6) placenta praevia;

7) congenital defects.

The correct answer is:

A. 1,4.

B. 2,6.

C. 2,7.

D. 3,5.

E. 4,6.

Pytanie nr **183**

The treatment of pre-invasive cervical cancer consists in:

- A.** excision of the affected part of the cervix.
- B.** strict cytological and colposcopic examination once a month.
- C.** simple hysterectomy.
- D.** radical hysterectomy.
- E.** brachytherapy.

Pytanie nr **184**

Which of the following ovarian tumors is a virilizing tumor?

- A.** Sertoli-Leydig cell tumor.
- B.** endometrioid tumor.
- C.** Wolffian duct cell tumor.
- D.** corpus luteum cyst.
- E.** fibroblast tumor.

Pytanie nr **185**

A woman who is rhesus negative undergoes spontaneous miscarriage at 14 week. What dose of anti D immunoglobulin should she receive?

- A.** 50 µg.
- B.** 100 µg.
- C.** 150 µg.
- D.** 300 µg.

E. 1500 µg.

Pytanie nr **186**

The relation of the long fetal axis to the long axis of the uterus and reproductive canal is called:

A. fetal lie.

B. position.

C. fetal location.

D. engagement.

E. presentation.

Pytanie nr **187**

A feature common for the Mayer-Rokitansky-Kustner-Hauser syndrome and complete androgen insensitivity syndrome is:

A. XY karyotype.

B. XX karyotype.

C. lack of the uterus and vagina.

D. defect of the androgen receptor gene.

E. presence of testicles in the abdominal cavity.

Pytanie nr **188**

When should the next cytological examination be ordered in a 35-year-old multipara with normal results of the last three cytological examinations and without any risk factors?

A. in 3 months.

B. in 6 months.

C. in a year.

D. in 3 years.

E. only if some symptoms occur, such as increased vaginal discharge, or contact bleeding.

Pytanie nr **189**

Indicate **the false** statement concerning hormonal contraception:

- A. progestogen-only pills can be used during breastfeeding.
- B. in the case of progestogen-only pills irregular bleedings or secondary amenorrhea can occur.
- C. hormonal contraception decreases the risk of ovarian, cervical and endometrial cancers.
- D. migraine is a contraindication to the use of oral hormonal contraception.
- E. so-called breakthrough bleedings are more intense in smokers.

Pytanie nr **190**

A patient with twin monochorionic pregnancy has been informed about the risk of twin-to-twin transfusion syndrome (TTTS). The syndrome is characterized by an unbalanced blood flow between the twins through active interconnecting blood vessels. One of the complications of TTTS that occurs in the recipient is:

- A. fetal hypertrophy.
- B. growth retardation.
- C. anemia.
- D. oligohydramnios.
- E. hypovolemia.

Pytanie nr **191**

Endometriosis should be taken into account in the case of:

- A. endometrial hyperplasia on ultrasound, the presence of symptoms of hirsutism.
- B. abnormal postmenopausal bleeding from the uterine cavity.
- C. reduced fertility, pelvic pain syndrome, dysmenorrhea.

D. rare menstruation with the image of polycystic ovaries on ultrasound.

E. presence of endometrial polyps in the uterine cavity.

Pytanie nr 192

The proliferative phase of the uterine cycle is dependent on:

A. progesterone produced by the corpus luteum.

B. progesterone produced by granulosa cells.

C. estrogen produced by the corpus luteum.

D. estrogen produced by granulosa cells.

E. estradiol produced by the dominant follicle.

Pytanie nr 193

Precocious puberty in girls is defined when tertiary sexual features appear before:

A. the age of 8.

B. the age of 8, but when menstruation starts concomitantly with pubic hair.

C. the age of 10.

D. the age of 10, but when menstruation starts concomitantly with pubic hair.

E. the age of 11 regardless of race.

Pytanie nr 194

Which of the following foetal head diameters is the smallest and most conveniently goes through the spaces of the pelvic planes during physiological childbirth?

A. suboccipitobregmatic.

B. occipitofrontal.

C. occipitomenal.

D. submentobregmatic.

E. verticomental.

Pytanie nr 195

Which of the following findings on cardiotocography (CTG) is a good prognostic factor?

A. decelerations.

B. sinusoidal pattern.

C. tachycardia.

D. accelerations.

E. bradycardia.

Pytanie nr 196

Which of the following changes in blood morphology occurs in the second trimester of normal pregnancy?

A. increased hematocrit levels.

B. decreased hemoglobin concentration.

C. increased hemoglobin concentration.

D. increased platelet count.

E. increased leucocyte count.

Pytanie nr 197

Directly after the third stage of the delivery massive hemorrhage occurred leading to hypovolemic shock. The condition of the patient improved significantly over several days, but she is suspected of Sheehan's syndrome. Which of the following justifies this suspicion?

A. her child receiving 7 in the Apgar score.

B. change in respiratory rate.

C. worsening of the general condition and depressive thoughts.

D. no lactation.

E. occurrence of diarrhea.

Pytanie nr **198**

Which of the following **is not** a physiological cause of increased prolactin secretion in women?

A. pregnancy.

B. strenuous activity.

C. sleeping.

D. stress.

E. increased glucose level.

Pytanie nr **199**

Low molecular weight heparin during pregnancy:

A. has tocolytic effect.

B. accelerates foetal lung maturation.

C. prevents miscarriage in women with hereditary thrombophilia.

D. decreases the risk of premature placental abruption.

E. should not be used due to its teratogenic effect.

Pytanie nr **200**

Indicate the true statement concerning Duncan's method of placental separation:

A. it is correlated with lower bleeding than in Schultze's method.

B. it begins centrally.

C. it occurs in 80% of cases.

D. more frequently occurs in twin pregnancy.

E. usually does not lead to a hematoma formation.

Pytanie nr **201**

The pharmacological treatment of endometriosis **does not** include:

- A. danazol.
- B. gonadoliberin analogues.
- C. preparations which contain 17-b estradiol only.
- D. progestogens.
- E. estrogen-progestogen formulations.

Pytanie nr **202**

Dopamine D₂ receptor agonists (bromocriptine, quinagolide, cabergoline) are used for:

- A. stimulation of lactation when the amount of milk produced in the postpartum period is too low.
- B. treatment of pituitary tumors producing prolactin.
- C. diagnostics of functional hyperprolactinemia.
- D. controlled ovarian hyperstimulation.
- E. contraception.

Pytanie nr **203**

Indicate the true statements concerning the cardiotocography (CTG) analysis:

- 1) short-term variability (STV) is particularly helpful in predicting the risk of metabolic acidosis in the fetus;
- 2) normal fetal heart rate (FHR) is 100 to 160 beats per minute;
- 3) lack of an acceleration in the record up to 180 minutes is regarded as normal and as a sign of deep sleep of the fetus;
- 4) correct long-term variability (LTV) is between 5 and 25 beats per minute;
- 5) sinusoidal rhythm indicates a high degree of fetal distress.

The correct answer is:

- A. 1,3,5.

B. 1,5.

C. 2,3.

D. 1,4,5.

E. 2,3,4.

Pytanie nr 204

The examiner places his hands flat and parallel on both sides of the abdomen of a pregnant woman, at the level of the umbilicus. The described maneuver allows to determine the fetal position and is called:

A. 1st Leopold's maneuver.

B. 2nd Leopold's maneuver.

C. 3rd Leopold's maneuver.

D. 4th Leopold's maneuver.

E. 5th Leopold's maneuver (Zangemeister maneuver).

Pytanie nr 205

Adaptive changes in the circulatory system of a pregnant woman consist of:

1) increase in the circulating blood volume, including 45% of the plasma volume and 20% of the erythrocyte mass, with a greater increase of the circulating blood volume in obese women than in slim women;

2) increase in cardiac output;

3) acceleration of heart rate by 10-15 beats per minute;

4) increased peripheral arterial resistance by 20%;

5) increase in central venous pressure.

The correct answer is:

A. 1,5.

B. 1,2,3.

C. 1,3,4.

D. only 1.

E. 4,5.

Pytanie nr **206**

A 27-year-old patient comes to the emergency department at 29 weeks of her first pregnancy complaining about a fever up to 38.3°C for the last couple of hours, 2 episodes of vomiting and back pain. Ultrasound examination reveals a normally growing living foetus, normal placenta located in the fundus and the full bladder. On palpation, the abdomen is not rigid but tender in the hypogastric area. On bimanual vaginal examination no pain during cervical motion is observed. The percussion of the lumbar area, especially on the left side, is very painful for the patient. What do you suspect?

- A. acute appendicitis.
- B. acute cystitis.
- C. left-sided renal calculi.
- D. acute pyelonephritis.
- E. hydronephrosis due to the pressure of the pregnant uterus on the ureter.

Pytanie nr **207**

All of the following are features of Swyer syndrome, **except from**:

- A. XY karyotype.
- B. uterus present.
- C. primary amenorrhea.
- D. male phenotype.
- E. hypogonadism.

Pytanie nr **208**

Which of the following are characteristic of a granuloma cell tumor?

- 1) most often occurs on both sides;
- 2) it is a germ cell tumor;
- 3) it is secreting estrogens and inhibin;
- 4) it is often accompanied by ascites;

5) most often occurs before puberty.

The correct answer is:

A. 1,2,4.

B. 2,4,5.

C. only 3.

D. 1,3,5.

E. 2,3,5.

Pytanie nr 209

In a pregnant (41 week of pregnancy) CTG test was done. It showed a basic heart rate 145/minute. The heart rate started to drop to 110/minute at the top of each contraction and it has been observed for the last 20 minutes. This record shows:

A. utero-placental insufficiency and it is an alarming sign because it precedes intrauterine foetal death.

B. foetal head compression and vagus nerve irritation and is not an adverse symptom unless it correlates with extended basic changes in the heart rate.

C. it is an indication for shortening the second stage of labour.

D. threat of intrauterine asphyxia and the need to consider a caesarean section.

E. none of the above.

Pytanie nr 210

Which of the history or diagnostics information **does not** confirm the diagnosis of hypothalamic-pituitary insufficiency?

A. amenorrhea.

B. high FSH levels.

C. low FSH levels.

D. negative progesterone test.

E. positive estrogen-progesterone test.

Pytanie nr **211**

Disulfiram-ethanol reaction may be expected in a patient with an infection of reproductive organs, who drank alcohol during the treatment with:

- A. metronidazole.
- B. cefuroxime.
- C. clindamycin.
- D. furaginum.
- E. acyclovir.

Pytanie nr **212**

Which of the following **should not** be used in the treatment of a urinary tract infection in a pregnant or breastfeeding woman?

- A. nitrofurantoin.
- B. ciprofloxacin.
- C. cefuroxime.
- D. amoxicillin.
- E. ampicillin.

Pytanie nr **213**

Which of the following is a non-invasive prenatal test that shows the highest sensitivity in the diagnostics of chromosomal aberrations in a fetus?

- A. chorionic villus sampling.
- B. amniocentesis.
- C. triple test.
- D. PAPP-A determination.

E. examination of cell-free fetal DNA in maternal blood.

Pytanie nr 214

Teratozoospermia according to WHO criteria of 2010 denotes:

A. percentage of sperm with normal morphology below 4%.

B. percentage of sperm showing normal mobility below 4%.

C. no sperm in the ejaculate.

D. over 250 million sperm in 1 mL of the ejaculate.

E. percentage of sperm with normal morphology below 40%.

Pytanie nr 215

Which of the following is characteristic of the vaginal discharge in a woman with bacterial vaginitis?

A. pH < 4.5, yellow-greenish foamy discharge with clue cells present.

B. pH < 4.5, grey or milky homogenous discharge with clue cells present.

C. pH > 4.5, grey or milky homogenous discharge with clue cells present.

D. pH > 4.5, white, clotty, serous discharge with clue cells present.

E. pH < 4.5, white, clotty, serous discharge without any cells present.

Pytanie nr 216

Indicate the true statement concerning ovarian cancer surgery:

A. it strives for tumour sample obtaining and fast radical chemotherapy application.

B. cytoreduction is performed to the highest possible extent.

C. only and always the affected ovary is removed.

D. following uterectomy the region of the lesser pelvis may be irradiated.

E. neoadjuvant chemotherapy is always used in order to decrease a tumour mass.

Pytanie nr **217**

Indicate the clinical features of germinal neoplasms of the ovary:

- 1) occur mainly in young women;
- 2) occur mainly in post-menopausal women;
- 3) may be diagnosed on the basis of increased blood levels of CA-125, CEA, CA 15-3;
- 4) may be diagnosed on the basis of increased blood levels of LDH, AFP, hCG;
- 5) respond well to chemotherapy;
- 6) are treated only surgically;
- 7) are always malignant tumours.

The correct answer is:

- A.** 1,4,7.
- B.** 3,4,6.
- C.** 2,4,6,7.
- D.** 1,4,5.
- E.** 2,3,7.

Pytanie nr **218**

Which of the following **is not** related to the polycystic ovary syndrome?

- A.** overweight and obesity.
- B.** insulin resistance and increased risk of type 2 diabetes.
- C.** increased risk of cardiovascular diseases.
- D.** increased risk of breast cancer and rectal cancer.
- E.** increased risk of endometrial cancer.

Pytanie nr **219**

Neoplastic ovarian tumours which produce steroid hormones include:

- 1) thecoma;
- 2) mature teratoma;
- 3) endometrioid carcinoma;
- 4) granulosa cell tumour.

The correct answer is:

A. 1,2.

B. 1,4.

C. 2,3.

D. 2,4.

E. 3,4.

Pytanie nr 220

The triad of Sabin and Pinkerton (hydrocephalus or microcephaly, chorioretinitis, intracranial calcifications) is typical of the congenital infection with:

A. rubella virus.

B. toxoplasmosis.

C. parvovirus B19.

D. cytomegalovirus.

E. herpes simplex virus.

Pytanie nr 221

The treatment of arterial hypertension in pregnant women includes the use of:

A. angiotensin-converting-enzyme inhibitors, methyldopa, labetalol.

B. angiotensin-converting-enzyme inhibitors, methyldopa, angiotensin II receptor antagonists.

C. labetalol, nifedipine, methyldopa.

D. diuretics, angiotensin II receptor antagonists, labetalol.

E. labetalol, methyldopa, amiodarone.

Pytanie nr 222

Which of the following **is not** a risk factor for endometrial cancer?

- A. smoking tobacco.
- B. obesity.
- C. diabetes.
- D. exogenous estradiol.
- E. late menopause.

Pytanie nr **223**

Which of the following can be usually found in a patient with menstrual disorders related to chronic intensive physical exercise?

- A. hypogonadotropic hypergonadism.
- B. hypergonadotropic hypergonadism.
- C. normal hormonal function of the ovaries.
- D. hypergonadotropic hypogonadism.
- E. hypogonadotropic hypogonadism.

Pytanie nr **224**

A **contraindication** to the safe delivery of the vaginal birth with the breech presentation of the fetus is:

- A. multiparity.
- B. pregnancy term 36-40 weeks.
- C. fetal frank breech presentation.
- D. fetal weight from 2000 to 3700 g.
- E. footlink breech presentation.

Pytanie nr **225**

The pregnancy in a patient with chronic hypertension is a high-risk pregnancy and may be complicated with the following disorders **except for**:

- A. preterm birth related to the cervical incompetence.
- B. intrauterine growth retardation.
- C. superimposed preeclampsia.
- D. abruption of normally located placenta.
- E. oligohydroamnios.

Pytanie nr **226**

Which of the following should be done in the case of a patient without risk factors for diabetes if her fasting glucose level is 110 mg/dL in the 9th week of gestation?

- A. oral glucose tolerance test with 75g of glucose between 24th and 28th week of gestation.
- B. repeated fasting glucose examination.
- C. recommend diabetes diet.
- D. assessment of the 24-hour blood glucose profile.
- E. oral glucose tolerance test with 75g of glucose immediately.

Pytanie nr **227**

Which of the following are **absolutely contraindicated** in the pharmacological treatment of arterial hypertension during pregnancy?

- A. calcium blockers.
- B. beta-blockers.
- C. thiazide diuretics.
- D. angiotensin convertase inhibitors.
- E. clonidine.

Pytanie nr **228**

Which of the following **is not** a typical complication of gestational diabetes?

- A. polyhydramnios.

- B. fetal macrosomia.
- C. intrauterine growth retardation.
- D. obstetric cholestasis.
- E. traumatic delivery.

Pytanie nr **229**

A **contraindication** to breastfeeding **is not**:

- A. HIV infection.
- B. breast cancer.
- C. severe mental disorder.
- D. mother's use of semisynthetic penicillin.
- E. acute heart failure.

Pytanie nr **230**

A young female patient gave birth to her child about a year ago through the vaginal delivery, after which curettage was necessary because of fragments of the placenta did not separate from the uterus. She stopped breastfeeding 4 months ago, but the periods have not reoccurred yet. Her medical history may suggest:

- A. Kallmann syndrome.
- B. Sheehan's syndrome.
- C. Asherman's syndrome.
- D. Mayer-Rokitansky-Küster-Hauser syndrome.
- E. Couvelaire syndrome.

Pytanie nr **231**

Which of the following blood vessels is used in the non-invasive ultrasound evaluation of foetal anaemia?

- A. uterine artery.
- B. ductus venosus.
- C. umbilical artery.
- D. middle cerebral artery.
- E. umbilical vein.

Pytanie nr **232**

High FSH serum levels accompanied by low estradiol levels are characteristic of:

- 1) Turner syndrome;
- 2) premature ovarian failure (POF);
- 3) pure gonadal dysgenesis;
- 4) polycystic ovary syndrome (PCOS);
- 5) postmenopausal period.

The correct answer is:

- A. 1,2,3.
- B. 1,2,5.
- C. 1,2,3,5.
- D. only 3.
- E. all the above.

Pytanie nr **233**

In obstetrics, the term position (*positio*) means:

- A. relationship of the foetus' body parts to the parts of the maternal womb.
- B. relationship of the longitudinal axis of the foetus to the longitudinal axis of the maternal womb.
- C. relationship of the foetal head to the foetal spine.
- D. relationship of the foetus' body parts to one another.
- E. relationship of a presenting part of the fetus to the inter-spinal line.

Pytanie nr **234**

Which of the following parameters **is not** included in the assessment of the fetal biophysical profile?

- A. fetal movement.
- B. fetal tone.
- C. umbilical artery blood flow.
- D. amniotic fluid volume.
- E. fetal heart rate.

Pytanie nr **235**

Indicate the true set of physiological changes in the circulatory system of pregnant women:

- A. increased heart rate, increased cardiac output, increased volume of circulating blood, decreased systemic and pulmonary vascular resistance, decreased arterial pressure.
- B. increased heart rate, decreased cardiac output, increased volume of circulating blood, increased systemic and pulmonary vascular resistance, decreased arterial pressure.
- C. increased heart rate, decreased cardiac output, increased volume of circulating blood, increased systemic and pulmonary vascular resistance, increased arterial pressure.
- D. increased heart rate, decreased cardiac output, decreased volume of circulating blood, increased systemic and pulmonary vascular resistance, decreased arterial pressure.
- E. decreased heart rate, decreased cardiac output, increased volume of circulating blood, increased systemic and pulmonary vascular resistance, increased arterial pressure.

Pytanie nr **236**

A 45-year-old woman with the diagnosis of cervical cancer went to see her gynecologist. Which of the following data from her medical history **is not** a risk factor for this cancer?

- A. 5 deliveries.
- B. numerous viral and bacterial infections of the cervix.

C. obesity.

D. smoking.

E. frequent changing of sexual partners.

Pytanie nr **237**

The complications typical of monochorionic-monoamniotic twin pregnancies include:

- 1) acute type of twin-to-twin transfusion syndrome (TTTS);
- 2) chronic type of twin-to-twin transfusion syndrome (TTTS);
- 3) impairment of the fetus whose twin died in the utero;
- 4) conjoined twins;
- 5) premature labor after the end of 37 week of gestation.

The correct answer is:

A. 2,3,4,5.

B. 1,3,4,5.

C. 1,2,3,4.

D. all the above.

E. 1,2.

Pytanie nr **238**

Anti-RhD immunoglobulin is used in RhD minus female patients in a single i.m. injection after the laparoscopic treatment of an ectopic pregnancy (WGA < 12 Hbd) within 72 h of the surgery at a dose of:

A. 50 µg.

B. 100 µg.

C. 150 µg.

D. 200 µg.

E. 300 µg.

Pytanie nr **239**

Which of the following **is not** included in the fetal biophysical profile (Manning's score)?

- A. fetal heart rate.
- B. fetal muscle tone.
- C. placenta development stage.
- D. amniotic fluid volume.
- E. fetal respiratory movements.

Pytanie nr **240**

The factors that increase the risk of endometrial cancer include:

- 1) smoking cigarettes;
- 2) metabolic syndrome (obesity, diabetes);
- 3) multiparity;
- 4) polycystic ovary syndrome;
- 5) long-term estrogen therapy not balanced with progesterone administration.

The correct answer is:

- A. 1,3,5.
- B. 3,4,5.
- C. 2,3,4.
- D. 2,4.
- E. 2,4,5.

Pytanie nr **241**

Which of the following fetal head circumferences is the smallest?

- A. occipito-frontal.
- B. suboccipito-bregmatic.
- C. occipito-mental.
- D. suboccipito-hyoid.

E. genio-hyoid.

Pytanie nr **242**

Premature *thelarche* in contrast to precocious puberty is characterized by the isolated:

A. development of axillary hair.

B. development of pubic hair.

C. breast development.

D. occurrence of ovulation.

E. occurrence of voice change.

Pytanie nr **243**

Marsupialization is a surgical method of:

A. primary treatment of stress urinary incontinence.

B. treatment of recurrent stress urinary incontinence.

C. treatment of a recurrent Bartholin gland abscess.

D. treatment of impaired statics of the reproductive organ.

E. creation of the vagina in patients with Mayer-Rokitansky-Küster-Hauser syndrome.

Pytanie nr **244**

A 32-year-old childless patient has a history of 4 miscarriages in the first trimester. The evaluation of the possible causes of miscarriages in this patient should include:

- 1) diagnostics of *Toxoplasma gondii* infection;
- 2) diagnostics of *Human Papilloma Virus* infection;
- 3) determination of anti-cardiolipin antibodies;
- 4) diagnostics of Cytomegalovirus infection;
- 5) determination of lupus anticoagulant.

The correct answer is:

A. 1,2.

B. 2,3.

C. 3,5.

D. 2,4,5.

E. 1,4,5.

Pytanie nr **245**

Which of the following are assessed by the Bishop score?

A. relation between the cervix and the vaginal axis, dilation, cervical effacement, fetal station.

B. gestational age, fetal position, cervical dilation and effacement.

C. cervical consistency, fetal lie, dilation, cervical effacement, fetal presentation.

D. relations between the uterine axis and vaginal axis, dilation, cervical effacement, fetal position.

E. relations between the long fetal axis and the vaginal axis, dilation, cervical effacement, strength of uterine muscle contractions.

Pytanie nr **246**

The diagnosis of endometriosis should be considered in all the following conditions, **except for**:

A. oligomenorrhea.

B. dyspareunia.

C. painful menstruation.

D. ovarian cyst with dense content.

E. pelvic pain.

Pytanie nr **247**

So called *clue cells* are typical of:

A. vaginal candidiasis.

- B.** vaginal trichomoniasis.
- C.** bacterial vaginosis.
- D.** normal vaginal biocenosis.
- E.** vaginal actinomycosis.

Pytanie nr **248**

The increased risk of endometrial cancer **has not** been associated with:

- A.** obesity.
- B.** diabetes.
- C.** hypertension.
- D.** late menopause.
- E.** endometriosis.

Pytanie nr **249**

Chemotherapy as a separate therapeutic strategy (not as a part of a combination therapy) is the most effective in the treatment of:

- A.** serous ovarian cancer.
- B.** mucinous ovarian cancer.
- C.** endometrial cancer of the uterus.
- D.** choriocarcinoma.
- E.** cervical squamous cell carcinoma.

Pytanie nr **250**

A 36-year-old multiparous woman went to see a gynecologist because of regular but excessive and painful periods which has lasted for the last few years. This is accompanied by increasing fatigue, general weakness, and more frequent urination. The pathological findings included pale mucosa. Gynecological exam revealed a tumour the size of a 4-month pregnancy. Blood test and transvaginal ultrasound were made. What lab test

results and diagnosis can you expect?

- A.** aneamia, uterine myomas.
- B.** anaemia, pregnancy.
- C.** normal blood count, pregnancy.
- D.** normal blood count, simple ovarian cyst.
- E.** anaemia, simple ovarian cyst.

Pytanie nr **251**

A 14-year-old girl went to see a gynecologist due to cramp pains within lower abdomen, accompanied by nausea, vomiting and general weakness. The pain started just before her menstruation and have continued for two days. A similar condition occurred during her last cycle. The girl is disturbed as she had her menarche the year before and did not have any similar symptoms. Gynecological exam did not reveal any abnormalities. Abdominal and pelvic ultrasound did not show any pathological findings. The most probable diagnosis will be:

- A.** premenstrual syndrome (PMS).
- B.** unicornate uterus.
- C.** primary dysmenorrhea.
- D.** secondary dysmenorrhea.
- E.** Asherman syndrome.

Pytanie nr **252**

Indicate which of the activities below is an example of primary prevention in gynecologic onology:

- A.** cytology.
- B.** HPV vaccination.
- C.** colposcopy.
- D.** HPV genotyping.

E. cervical conization/LLETZ.

Pytanie nr **253**

Influenza vaccination in pregnancy:

- 1) is recommended to every woman who has no contraindications to vaccination;
- 2) should be performed only in the second trimester of pregnancy;
- 3) should be performed using inactivated quadrivalent vaccine;
- 4) is safe both for the mother and the child.

The correct answer is:

- A. 1,4.
- B. all of the above.
- C. 1,3,4.
- D. 1,2,4.
- E. 1,2.

Pytanie nr **254**

The Mayer-Rokitansky-Kustner and Hauser syndrome is characterized by:

- 1) primary amenorrhoea;
- 2) female karyotype 46,XX;
- 3) scarcity of axillary and pubic hair;
- 4) congenital absence of ovary;
- 5) congenital absence of vagina;
- 6) ametria, rudimentary uterus or rudimentary horns of uterus.

The correct answer is:

- A. all of the above.
- B. 1,2,5,6.
- C. 2,3,4,5.
- D. 2,3,4,6.
- E. 1,4,5,6.

Pytanie nr **255**

A patient in a 36-week pregnancy had contact with a person diagnosed with chickenpox. If the patient is non-immune to chickenpox, the correct proceeding will be:

- A.** immunoglobulin given only to the newborn, just after birth.
- B.** informing the patient that no prophylactic treatment is necessary.
- C.** intramuscular immunoglobulin given to the patient, within 72-96 hours from contact.
- D.** intramuscular immunoglobulin given to the patient, within 72-96 hours from contact, and to the newborn, immediately after birth.
- E.** performing a cesarean section within 24 hours from contact with the chickenpox-infected person, and immunoglobulin given to the newborn just after birth.

Pytanie nr **256**

In the case of post-menopausal bleeding, the recommended diagnostic procedure will be:

- 1) ultrasound examination after 2 months;
- 2) endometrial aspiration biopsy;
- 3) hysteroscopy and targeted biopsy;
- 4) computed tomography (CT) of pelvis;
- 5) fractional curettage of the uterine cavity.

The correct answer is:

- A.** 1,2,3.
- B.** 1,2,5.
- C.** 1,2,3,5.
- D.** 4 only.
- E.** 2,3,5.

Pytanie nr **257**

A 28-year-old patient obtained an ASC-H cytology result. The patient was not pregnant and was not vaccinated against HPV. The recommended procedure will be:

- A.** colposcopy.
- B.** cervical conization, both for diagnostic and therapeutic purposes.

C. HPV genotyping.

D. assessment of the p16/Ki67 biomarkers.

E. repeated cytology after 6 months.

Pytanie nr **258**

The risk factors for gestational diabetes **do not** include:

A. obesity of pregnant woman.

B. diabetes in first-degree relatives.

C. diabetes in previous pregnancy.

D. age above 35 years.

E. diabetic father of the child.

Pytanie nr **259**

Oligohydramnios can be caused by:

1) fetal lung hypoplasia;

2) premature rupture of membranes;

3) fetal hypotrophy;

4) placental insufficiency;

5) viral infection.

The correct answer is:

A. 1,2,3.

B. 2,4,5.

C. 2,3,4.

D. 3,4,5.

E. 1,4,5.

Pytanie nr **260**

Endometriosis is a disease in which endometrial glands and stroma occur outside the

uterus. The clinical symptoms of endometriosis include:

- A.** dyspareunia.
- B.** infertility.
- C.** dysmenorrhea.
- D.** pelvic pain syndrome.
- E.** all of the above.

Pytanie nr 261

A 60-year-old female visited her gynaecologist because of vaginal bleeding/spotting. The symptoms appeared after sexual intercourse. The patient has not had periods for 10 years. She has not observed any spotting or bleeding so far. For the last few years she has been treated with marcumar for atrial fibrillation. Indicate the possible cause(s) of these symptoms:

- A.** the use of marcumar.
- B.** cervical cancer.
- C.** endometrial proliferation/cancer.
- D.** vaginal atrophy.
- E.** all of the above.

Pytanie nr 262

A 14-year-old girl was found to have no breast growth. This may be the symptom of:

- 1) Rokitansky syndrome;
- 2) gonadal dysgenesis;
- 3) constitutional delay of puberty;
- 4) androgen insensitivity syndrome;
- 5) some metabolic diseases, e.g. diabetes.

The correct answer is:

- A.** 1,2,3.
- B.** 2,3,4.

C. 3,4,5.

D. 2,3,5.

E. all of the above.

Pytanie nr **263**

In imminent premature labour, the **absolute** contraindications to tocolytic treatment **do not** include:

A. fetal death.

B. intrauterine infection.

C. premature rupture of membranes.

D. premature placental abruption.

E. eclampsia.

Pytanie nr **264**

In week 27 of diamniotic monochorionic twin pregnancy, one of the foetuses was found to have oligohydramnios, collapsed bladder, and reduced fetal movement, while the other had overfull bladder and generalised oedema. The diagnosis should be:

A. selective intrauterine growth restriction (sIUGR), and the treatment should be medical observation and another ultrasound after two weeks.

B. reversed flow syndrome, and the treatment should be a sclerosant injected into the artery of the acardiac foetus.

C. twin-to-twin transfusion syndrome (TTTS), and the treatment should be fetoscopic lasercoagulation of blood vessels connections between the twins.

D. twin anaemia-polycythemia sequence, and the treatment should be urgent termination of pregnancy via the caesarean section.

E. comorbidity of gestational trophoblastic disease, and the treatment should be urgent termination of pregnancy via the caesarean section.

Pytanie nr **265**

In diabetic patients, the contraindications to pregnancy **do not** include:

- A. severe nephropathy.
- B. treatment-resistant arterial hypertension.
- C. ischaemic heart disease.
- D. diabetic nephropathy.
- E. treatment-resistant proliferative retinopathy.

Pytanie nr **266**

Indicate true statements regarding epidemiology of cervical cancer:

- A. the incidence of cervical cancer is correlated with the rate of human papilloma virus.
- B. the relative risk of cervical cancer decreases with age.
- C. cervical cancer is more prevalent among obese women.
- D. cervical cancer is not found among sexually inactive women.
- E. the relative risk of cervical cancer decreases with the number of childbirths.

Pytanie nr **267**

The most frequent type of vulvar cancer (65-90% of cases) is:

- A. adenocarcinoma.
- B. squamous cell carcinoma.
- C. clear cell carcinoma.
- D. Paget's disease.
- E. Bowen's disease.

Pytanie nr **268**

A female genital neoplasm which gives a positive pregnancy test result, and is often manifested by the development of bilateral ovarian cysts during its course, is termed as:

- A. *dysgerminoma*.
- B. *folliculoma*.
- C. *sertolioma*.
- D. *choriocarcinoma*.
- E. *luteoma*.

Pytanie nr **269**

In gynaecology, an example of primary prevention of cervical cancer is:

- A. DNA genotyping of highly oncogenic HPVs.
- B. cytology.
- C. a so-called 'co-testing' i.e. cytology combined with HVP genotyping.
- D. vaccination against HPV.
- E. colonoscopy.

Pytanie nr **270**

The basic examination for detecting cervical cancer is:

- A. routine gynaecological examination.
- B. cytological assessment of the peritoneal fluid obtained in abdominal paracentesis.
- C. determination of Ca125 marker.
- D. ROMA test (Ca125 and HE-4 markers).
- E. transvaginal ultrasound of pelvic organs.

Pytanie nr **271**

In ovarian cancer diagnostics, the family history of this cancer is of high significance. Which genes should be considered in gene mutation testing that could help assess an increased risk of ovarian cancer among family members?

- A. *P 53*.

B. Bcl-2.

C. BRCA1 and BRCA2.

D. ACE 2.

E. HE4.

Pytanie nr **272**

Indicate which of the factors below contribute to decreasing the risk of ovarian cancer and are thought to play a protective role:

- 1) multiparity;
- 2) combined hormonal contraceptive;
- 3) hormonal stimulation of ovulation;
- 4) ovarian puncture in the *in vitro* fertilization procedure.

The correct answer is:

A. 2 only.

B. 1,2.

C. 1,2,3.

D. all of the above.

E. none of the above.

Pytanie nr **273**

Indicate hormonal drugs used in the complementary therapy of cancer of the body uterus:

A. estrogens.

B. selective oestrogen receptor modulators.

C. progestagens.

D. selective progesterone receptor modulators.

E. androgens.

